

קידום בריאות																
מאחדת				מכבי				לאומית				כלית				← SES ↓ קבוצת גיל ↓ מין
8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	
שיעור התעודות של מרכיבי BMI בקרב בני 64-20																
60.9%	57.1%	59.7%	60.5%	63.5%	64.5%	60.2%	60.0%	54.9%	53.8%	51.5%	49.4%	72.0%	72.8%	70.1%	72.9%	20-24
76.0%	77.0%	78.1%	79.8%	59.0%	63.0%	63.2%	60.9%	71.1%	73.3%	72.2%	71.4%	71.4%	72.8%	70.1%	68.4%	25-34
84.0%	84.0%	83.6%	85.3%	63.7%	68.2%	68.5%	68.1%	76.2%	76.5%	76.5%	75.9%	78.6%	79.2%	78.1%	78.2%	35-44
92.4%	91.8%	90.9%	92.0%	71.9%	76.7%	77.4%	78.7%	84.5%	84.3%	84.1%	85.0%	86.6%	88.1%	88.1%	88.6%	45-54
95.4%	95.2%	94.5%	95.2%	81.1%	85.2%	86.1%	87.4%	91.8%	91.1%	90.4%	90.0%	95.0%	96.1%	96.0%	96.5%	55-64
74.4%	73.4%	77.4%	80.2%	72.3%	77.9%	79.9%	81.2%	67.3%	67.3%	67.9%	68.1%	80.7%	81.6%	81.1%	82.9%	20-24
88.9%	91.0%	93.0%	95.6%	78.4%	83.7%	85.3%	86.7%	88.6%	90.3%	91.3%	92.3%	81.2%	84.0%	83.5%	82.3%	25-34
91.6%	91.8%	93.9%	95.3%	75.1%	79.8%	81.1%	83.0%	86.8%	88.2%	89.1%	91.3%	84.1%	86.0%	86.6%	86.9%	35-44
94.0%	94.1%	94.3%	95.8%	74.4%	80.4%	82.2%	84.0%	88.2%	89.2%	90.0%	89.8%	89.1%	91.7%	92.1%	92.7%	45-54
96.0%	96.3%	95.7%	96.9%	81.8%	86.1%	87.9%	89.1%	92.9%	93.8%	93.6%	93.2%	96.3%	97.1%	97.4%	98.0%	55-64
שיעור התמנאות של תת משקל בקרב בני 64-20																
6.9%	7.2%	6.0%	5.8%	7.6%	7.0%	7.1%	5.8%	6.2%	3.7%	5.1%	5.9%	8.2%	8.3%	5.8%	5.2%	20-24
2.9%	2.9%	3.4%	3.0%	2.4%	2.7%	3.3%	3.7%	2.9%	3.0%	3.6%	3.6%	2.4%	3.2%	3.6%	3.0%	25-34
1.4%	1.4%	1.9%	1.9%	1.0%	1.2%	1.5%	1.6%	1.7%	1.7%	1.8%	1.9%	1.2%	1.6%	1.9%	1.7%	35-44
0.7%	0.7%	1.1%	1.0%	0.4%	0.5%	0.9%	0.7%	0.9%	0.9%	1.1%	0.9%	0.6%	0.9%	1.2%	1.1%	45-54
0.5%	0.6%	0.9%	0.8%	0.3%	0.4%	0.7%	1.1%	0.6%	0.8%	0.9%	1.2%	0.5%	0.8%	1.3%	1.4%	55-64
10.0%	9.3%	7.4%	6.7%	11.6%	9.8%	8.5%	7.3%	9.3%	9.6%	8.9%	6.8%	12.0%	11.8%	11.1%	8.6%	20-24
6.2%	5.6%	4.4%	3.3%	6.0%	5.0%	4.5%	3.4%	7.8%	5.3%	4.4%	3.8%	7.3%	6.5%	6.4%	4.2%	25-34
4.4%	3.5%	3.0%	1.9%	4.0%	3.0%	2.7%	1.8%	4.5%	3.5%	2.7%	1.8%	5.1%	4.2%	3.4%	1.7%	35-44
2.6%	2.2%	1.7%	1.1%	2.9%	2.0%	1.7%	1.1%	2.8%	2.0%	1.7%	1.1%	3.1%	2.5%	1.9%	1.0%	45-54
1.6%	1.4%	1.2%	0.8%	1.9%	1.4%	1.3%	0.9%	2.1%	1.6%	1.2%	0.9%	1.8%	1.4%	1.3%	0.9%	55-64
שיעור התמנאות של עודף משקל בקרב בני 64-20																
25.2%	24.3%	25.1%	25.7%	20.9%	22.2%	23.7%	24.9%	25.3%	25.9%	28.5%	26.5%	21.9%	22.2%	24.8%	22.7%	20-24
29.2%	32.1%	33.2%	32.9%	29.2%	31.6%	31.5%	32.2%	26.9%	32.0%	32.9%	33.1%	29.0%	31.1%	31.7%	32.0%	25-34
37.9%	39.2%	38.7%	37.9%	37.8%	39.1%	37.6%	36.5%	37.4%	39.1%	39.5%	37.9%	37.2%	38.4%	38.0%	38.9%	35-44
44.3%	43.2%	41.5%	39.6%	44.8%	43.8%	41.3%	39.3%	44.3%	42.7%	41.9%	41.4%	43.6%	42.6%	41.6%	42.4%	45-54
48.1%	46.2%	42.7%	41.0%	47.9%	46.3%	43.3%	41.5%	47.3%	44.9%	42.8%	41.6%	48.0%	44.6%	42.9%	41.9%	55-64
21.7%	23.2%	25.1%	27.7%	18.4%	21.5%	25.2%	27.7%	16.8%	21.9%	24.8%	27.2%	16.2%	18.7%	20.3%	21.2%	20-24
26.0%	28.5%	31.0%	32.5%	27.1%	29.7%	30.4%	33.0%	25.3%	29.4%	29.7%	32.7%	21.9%	24.6%	25.9%	28.9%	25-34
26.7%	29.4%	31.8%	34.0%	29.9%	30.9%	31.2%	34.0%	26.9%	29.7%	33.0%	33.9%	24.8%	27.1%	29.7%	33.1%	35-44
29.3%	31.3%	33.4%	33.7%	28.3%	31.6%	32.3%	32.4%	29.7%	31.6%	33.8%	33.2%	27.8%	30.3%	32.6%	33.4%	45-54
34.3%	35.3%	35.0%	31.7%	33.1%	35.0%	34.9%	32.2%	32.9%	35.3%	34.4%	32.5%	33.6%	34.7%	33.8%	31.3%	55-64
שיעור ההתמנאות של השמנת יתר בקרב בני 64-20																
13.6%	14.3%	16.0%	17.3%	12.3%	16.6%	16.8%	17.5%	11.6%	14.7%	15.5%	17.9%	12.4%	15.5%	15.1%	14.0%	20-24
11.2%	15.6%	19.9%	23.0%	11.8%	16.9%	20.7%	23.4%	11.4%	17.2%	19.1%	22.5%	11.5%	17.1%	18.1%	20.4%	25-34
14.9%	22.1%	26.8%	32.2%	16.2%	22.4%	27.9%	33.4%	16.7%	22.2%	25.4%	31.1%	14.3%	19.4%	23.5%	26.1%	35-44
21.5%	28.2%	32.6%	37.6%	21.8%	29.5%	34.2%	39.1%	22.7%	29.3%	32.5%	35.8%	19.6%	25.5%	29.1%	31.1%	45-54
23.7%	29.3%	33.9%	35.9%	24.8%	30.0%	33.7%	36.5%	25.4%	30.5%	31.2%	34.3%	24.0%	28.4%	29.1%	30.7%	55-64
13.5%	15.1%	16.3%	18.8%	10.9%	16.3%	17.6%	17.9%	12.2%	16.1%	18.0%	20.0%	11.8%	14.4%	14.6%	15.6%	20-24
16.5%	21.7%	24.8%	27.5%	15.5%	22.9%	26.9%	28.5%	16.5%	24.0%	26.3%	29.0%	13.3%	19.4%	22.1%	25.8%	25-34
18.2%	24.2%	30.1%	36.9%	19.0%	25.4%	30.0%	36.0%	20.6%	26.2%	29.8%	39.1%	15.8%	22.1%	27.2%	34.4%	35-44
19.0%	26.1%	33.9%	44.7%	19.8%	27.5%	34.7%	43.3%	21.5%	29.5%	36.1%	46.2%	19.5%	26.6%	34.5%	44.9%	45-54
22.1%	29.1%	38.8%	51.5%	21.9%	30.9%	37.6%	47.8%	25.2%	31.6%	41.4%	49.0%	23.5%	31.2%	39.8%	51.1%	55-64
שיעור היעדר תעודות מצב שישן בקרב בני 74-16																
55.7%	51.8%	50.7%	58.2%	64.4%	69.3%	71.8%	71.6%	71.9%	72.3%	73.7%	71.8%	98.2%	99.1%	99.2%	99.6%	16-24
59.7%	57.5%	56.0%	62.1%	70.7%	74.4%	75.3%	76.5%	71.0%	71.1%	72.0%	71.8%	98.6%	99.2%	99.3%	99.7%	25-34
66.7%	63.7%	62.0%	67.0%	55.0%	60.8%	63.7%	63.3%	73.6%	73.9%	74.3%	75.6%	91.3%	93.3%	94.2%	94.4%	35-44
74.5%	71.9%	72.8%	77.3%	59.6%	67.5%	69.7%	70.4%	80.1%	81.1%	80.3%	83.8%	57.8%	63.3%	66.3%	66.7%	45-54
82.6%	81.2%	82.4%	85.4%	71.5%	77.7%	80.2%	81.4%	89.6%	88.4%	87.3%	88.8%	64.5%	72.0%	75.1%	75.9%	55-64
93.6%	92.0%	90.8%	91.0%	86.6%	89.0%	89.7%	89.7%	94.9%	93.3%	91.3%	91.6%	73.9%	76.8%	79.2%	79.6%	65-74
61.9%	58.2%	59.6%	68.5%	75.7%	81.2%	81.2%	80.8%	80.3%	81.2%	81.4%	80.7%	98.5%	99.3%	99.6%	99.8%	16-24
82.4%	83.2%	85.9%	89.7%	91.6%	93.0%	93.1%	94.1%	87.0%	89.5%	90.2%	91.8%	99.3%	99.7%	99.8%	99.9%	25-34
81.5%	78.6%	82.2%	87.1%	70.3%	72.5%	74.5%	74.3%	86.9%	87.5%	87.9%	90.9%	91.7%	92.8%	92.3%	90.1%	35-44
79.1%	76.2%	78.9%	84.5%	63.2%	69.7%	72.1%	71.8%	87.7%	87.8%	87.4%	89.4%	52.6%	53.0%	46.1%	31.0%	45-54
84.1%	82.5%	84.4%	88.3%	72.6%	77.1%	79.7%	80.0%	91.6%	91.8%	90.5%	91.6%	59.2%	59.6%	56.9%	44.2%	55-64
93.8%	91.5%	91.2%	92.2%	85.5%	87.4%	88.5%	88.7%	95.1%	94.3%	92.6%	93.1%	66.7%	65.1%	63.3%	51.8%	65-74
שיעור היעדר תעודות מצב שישן בקרב בני 74-16																
10.3%	15.6%	20.4%	31.4%	7.2%	11.2%	15.5%	16.8%	19.7%	23.2%	29.3%	29.6%	6.6%	11.7%	14.9%	15.1%	16-24
30.9%	37.0%	40.9%	51.5%	22.2%	27.5%	30.5%	27.2%	37.5%	42.6%	48.5%	49.7%	24.7%	31.7%	35.9%	38.1%	25-34
26.1%	32.8%	35.9%	45.4%	21.5%	28.6%	33.5%	30.8%	35.6%	38.8%	44.5%	48.4%	21.7%	30.4%	36.3%	42.2%	35-44
20.0%	30.3%	34.9%	44.3%	16.5%	24.9%	32.5%	30.6%	28.7%	37.8%	43.8%	46.3%	26.9%	39.6%	47.7%	56.1%	45-54
18.9%	28.3%	34.1%	42.2%	13.2%	22.0%	29.7%	29.7%	26.1%	34.8%	40.4%	42.6%	21.2%	34.2%	42.8%	51.7%	55-64
14.7%	20.5%	23.8%	32.4%	10.0%	15.7%	20.5%	21.0%	20.2%	25.6%	31.4%	33.8%	13.4%	21.4%	29.0%	38.8%	65-74
6.8%	7.8%	4.1%	4.6%	6.4%	6.8%	5.3%	1.9%	12.2%	12.9%	7.9%	5.2%	5.0%	5.7%	4.0%	1.3%	16-24
17.9%	17.8%	8.8%	7.4%	11.1%	12.3%	11.4%	5.4%	24.1%	23.0%	14.5%	10.0%	13.4%	15.3%	12.4%	4.6%	25-34
16.4%	19.5%	11.5%	8.5%	11.1%	15.3%	16.2%	8.5%	22.6%	24.6%	16.7%	11.2%	13.3%	17.9%	15.1%	6.9%	35-44
16.0%	20.1%	15.4%	10.5%	12.2%	17.7%	19.4%	11.5%	21.8%	25.1%	18.2%	12.3%	21.7%	31.0%	30.7%	24.2%	45-54
15.5%	20.5%	17.1%	13.0%	12.1%	16.5%	19.0%	12.8%	21.2%	24.3%	19.8%	16.8%	20.0%	28.3%	28.3%	26.2%	55-64
12.1%	14.9%	13.4%	9.6%	9.7%	12.3%	12.1%	9.3%	17.8%	19.4%	16.8%	14.4%	13.3%	18.5%	19.7%	18.9%	65-74

חא מסומן ב- * מייצג ערך מכנה נומרן 30 מבוססים

בדיקות סקר לגילוי מוקדם של סרטן																
מאוחדת				מכבי				לאומית				כללית				← SES קבוצת גיל ↓ מין ↓
8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	
שיעור הביצוע של ממוגנפיה לגילוי מוקדם של סרטן השד בנשים בנות 74-50																
73.6%	71.3%	65.0%	57.9%	79.6%	78.5%	75.1%	67.9%	64.9%	59.9%	56.1%	50.9%	75.1%	72.1%	68.9%	67.0%	50-54
71.1%	69.4%	63.3%	57.4%	79.0%	77.9%	74.4%	68.6%	58.8%	58.3%	54.3%	49.8%	73.4%	71.2%	68.9%	66.8%	55-59
70.8%	69.0%	64.4%	55.8%	78.3%	78.0%	75.6%	70.1%	62.4%	60.7%	55.1%	51.0%	74.6%	72.5%	70.1%	68.1%	60-64
70.8%	69.8%	65.2%	56.0%	77.8%	78.2%	75.6%	71.7%	61.5%	60.2%	56.8%	49.2%	75.8%	73.9%	70.2%	67.3%	65-69
69.5%	68.0%	61.8%	50.9%	74.1%	72.6%	70.8%	66.1%	59.5%	59.0%	53.2%	49.8%	71.3%	67.6%	62.5%	55.5%	70-74
שיעור הנשים בגילאים 54-25 אשר ביצעו בדיקות סיקור לסרטן צוואר הרחם בתדירות המומלצת																
62.2%	57.5%	46.0%	39.0%	77.6%	75.5%	69.1%	56.2%	71.6%	64.8%	56.4%	45.7%	74.7%	69.0%	58.5%	42.9%	25-29
70.4%	64.4%	54.3%	47.5%	88.8%	87.5%	81.5%	70.7%	75.5%	71.4%	60.9%	53.1%	84.1%	79.0%	69.8%	54.2%	30-34
67.6%	63.9%	52.7%	46.7%	88.2%	87.3%	82.3%	72.4%	75.6%	72.0%	62.1%	53.8%	83.4%	78.4%	69.0%	55.1%	35-39
68.8%	63.0%	52.5%	44.1%	87.1%	85.5%	80.3%	72.5%	74.4%	70.5%	59.6%	51.2%	79.9%	76.3%	66.0%	52.2%	40-44
67.1%	61.8%	51.9%	42.2%	85.3%	83.8%	78.7%	72.4%	71.2%	68.6%	59.0%	49.4%	77.4%	72.2%	62.1%	50.7%	45-49
63.0%	57.1%	47.3%	37.3%	83.4%	81.4%	76.4%	69.8%	68.1%	63.9%	54.4%	45.4%	72.7%	66.6%	55.4%	44.4%	50-54
שיעור הביצוע של בדיקה לגילוי מוקדם של סרטן המעי הגס בקרב בני 74-50																
56.0%	49.7%	45.0%	43.9%	62.9%	58.8%	53.7%	51.7%	49.9%	44.6%	41.2%	37.2%	55.5%	50.4%	47.5%	49.5%	50-54
66.2%	60.4%	53.5%	52.7%	73.5%	68.8%	63.7%	61.6%	59.8%	54.2%	49.4%	44.1%	65.5%	58.7%	55.9%	56.4%	55-59
68.3%	65.5%	60.6%	57.7%	76.4%	73.4%	69.1%	66.0%	65.9%	59.6%	54.0%	49.5%	68.8%	64.0%	60.0%	61.4%	60-64
73.0%	70.7%	65.6%	63.1%	78.8%	76.4%	73.8%	72.3%	66.3%	65.4%	61.2%	52.9%	73.4%	69.9%	65.5%	65.8%	65-69
76.5%	72.6%	69.2%	66.8%	81.0%	78.6%	75.9%	71.5%	71.9%	69.7%	63.6%	59.4%	76.2%	71.4%	67.4%	66.3%	70-74
52.6%	52.0%	48.9%	48.2%	61.2%	60.3%	57.7%	55.5%	50.7%	50.2%	46.4%	41.2%	54.4%	52.8%	52.8%	56.9%	50-54
63.7%	60.9%	56.2%	56.4%	70.4%	69.0%	66.0%	63.3%	57.3%	56.8%	53.5%	47.3%	64.5%	59.9%	59.3%	63.4%	55-59
68.1%	65.9%	61.6%	61.1%	74.4%	72.9%	71.5%	69.1%	63.5%	62.3%	57.3%	53.0%	68.3%	64.9%	63.7%	66.2%	60-64
71.5%	69.6%	66.9%	64.7%	77.5%	77.1%	74.5%	72.0%	68.0%	66.9%	62.6%	53.6%	72.5%	69.7%	67.1%	68.1%	65-69
71.9%	72.7%	67.4%	62.7%	77.4%	76.9%	74.7%	72.6%	68.9%	69.9%	65.0%	61.0%	72.9%	70.5%	66.6%	64.4%	70-74
שיעור המבוססים בעלי בדיקת דם סמי חיובית, אשר עברו בדיקת קולונסקופיה תוך 180 ימים מיום ביצוע בדיקת הדם הסמי																
68.3%	62.2%	51.6%	57.6%	84.9%	82.0%	80.3%	78.8%	NA*	51.1%	57.9%	55.6%	77.0%	69.6%	61.0%	53.5%	50-54
55.6%	68.8%	58.1%	44.1%	86.7%	80.3%	70.1%	71.1%	NA*	53.5%	42.1%	45.3%	73.5%	68.3%	58.4%	51.6%	55-59
NA*	54.4%	60.0%	53.8%	77.8%	77.0%	71.8%	56.5%	NA*	53.1%	48.2%	40.9%	70.1%	68.4%	58.9%	53.4%	60-64
56.8%	67.0%	52.9%	56.2%	78.9%	76.0%	67.0%	67.5%	NA*	61.3%	56.9%	40.4%	75.4%	68.6%	58.0%	49.0%	65-69
66.7%	56.8%	60.6%	52.9%	69.6%	70.6%	66.4%	58.5%	NA*	56.1%	45.1%	28.9%	75.6%	67.4%	58.3%	54.1%	70-74
51.6%	70.0%	63.0%	59.1%	83.3%	85.1%	81.9%	72.0%	NA*	54.3%	55.8%	42.2%	77.3%	70.4%	66.6%	59.9%	50-54
NA*	66.0%	54.3%	61.5%	83.5%	81.9%	72.8%	82.1%	NA*	63.4%	62.3%	60.0%	71.6%	73.7%	63.8%	53.8%	55-59
67.4%	61.5%	57.7%	65.6%	81.6%	76.6%	78.9%	70.0%	NA*	45.3%	65.8%	51.8%	74.5%	69.8%	65.7%	50.4%	60-64
69.7%	64.6%	66.7%	49.3%	81.2%	73.6%	73.8%	63.8%	NA*	57.3%	68.6%	35.0%	76.2%	70.3%	63.8%	50.6%	65-69
67.4%	64.0%	52.3%	53.3%	75.5%	73.8%	67.9%	68.4%	NA*	61.4%	52.7%	42.9%	76.4%	69.8%	59.0%	50.6%	70-74

חא מסומן ב-* מייצג ערך מכנה נמוך מן 30 מבוססים

ילדים ובני נוער																	
מאוחדת				מכבי				לאומית				כללית				← SES	
8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	מין ↓	
שיעור הביצוע של בדיקת המוגלובין בתוקף בני 18-9 חודשים																	
86.0%	84.9%	84.5%	85.4%	84.7%	81.6%	75.3%	76.2%	77.7%	74.0%	69.3%	68.5%	88.8%	86.7%	85.1%	88.6%	בנים	
85.5%	83.6%	83.7%	85.3%	83.1%	81.2%	73.8%	76.1%	80.2%	70.8%	66.1%	67.3%	87.6%	86.5%	84.1%	87.2%	בנות	
שיעור ההמצאות של אנמיה בקרב תינוקות בני 18-9 חודשים																	
7.5%	8.4%	12.3%	12.1%	9.0%	8.4%	10.5%	10.5%	9.9%	7.8%	11.8%	12.9%	9.3%	9.5%	9.2%	8.8%	בנים	
7.1%	7.9%	10.8%	11.3%	8.4%	8.4%	10.2%	11.1%	8.8%	7.7%	8.8%	10.2%	8.3%	8.8%	9.2%	8.7%	בנות	
שיעור התיעוד של מדידת טבע ומשקל בקרב ילדים בני 7																	
74.8%	70.2%	67.9%	71.3%	53.3%	53.6%	50.1%	47.9%	74.8%	72.3%	63.0%	59.1%	77.8%	75.5%	70.2%	70.5%	בנים	
72.6%	68.2%	69.1%	71.4%	53.3%	51.5%	48.6%	46.4%	69.7%	71.1%	63.6%	55.7%	78.3%	75.4%	69.1%	69.2%	בנות	
שיעור ההמצאות של רזון (Thinness) בקרב ילדים שמלאו להם 7 שנים בשנת המדד																	
4.6%	4.9%	5.7%	5.9%	4.1%	3.3%	4.1%	4.5%	3.8%	4.0%	4.3%	4.5%	3.8%	4.7%	5.2%	7.2%	בנים	
5.0%	4.1%	4.7%	5.0%	1.7%	2.9%	2.1%	2.6%	3.5%	3.2%	2.5%	3.4%	2.9%	3.2%	3.6%	6.6%	בנות	
שיעור ההמצאות של עודף משקל בקרב ילדים שמלאו להם 7 שנים בשנת המדד																	
14.8%	14.2%	12.6%	11.6%	10.6%	11.4%	9.9%	10.5%	10.1%	10.7%	12.9%	10.6%	10.4%	11.0%	10.6%	10.3%	בנים	
16.4%	14.2%	13.3%	12.1%	13.3%	13.3%	12.5%	11.8%	13.5%	13.5%	10.8%	11.7%	11.2%	12.5%	12.3%	11.0%	בנות	
שיעור ההמצאות של השמנת יתר בקרב ילדים שמלאו להם 7 שנים בשנת המדד																	
5.5%	8.9%	7.6%	8.8%	7.5%	8.4%	8.9%	8.4%	10.8%	9.7%	8.1%	8.3%	6.3%	8.3%	9.3%	9.3%	בנים	
6.6%	8.1%	7.4%	6.9%	6.9%	8.9%	10.0%	7.1%	9.6%	8.0%	9.8%	7.3%	5.6%	8.1%	9.3%	8.7%	בנות	
שיעור התיעוד של מרכיבי BMI בקרב בני 15-14																	
80.6%	73.5%	75.3%	75.8%	62.8%	62.7%	59.5%	56.6%	83.5%	81.2%	76.4%	67.3%	77.2%	76.0%	71.5%	72.4%	14	בנים
84.0%	79.8%	77.5%	79.3%	68.3%	68.0%	63.9%	56.3%	81.4%	74.1%	76.7%	67.3%	77.7%	76.3%	70.0%	60.9%	15	בנים
77.2%	70.2%	72.6%	74.8%	60.9%	61.0%	58.2%	55.0%	80.0%	79.3%	73.3%	64.8%	76.5%	76.1%	72.1%	71.9%	14	בנות
84.9%	77.5%	74.9%	80.1%	66.6%	67.1%	60.5%	54.8%	67.6%	81.3%	70.8%	63.0%	77.4%	76.3%	67.9%	63.3%	15	בנות
שיעור ההמצאות של רזון (Thinness) בקרב בני 15-14																	
4.0%	3.6%	4.6%	3.9%	4.0%	4.8%	4.5%	5.1%	3.8%	4.6%	4.8%	5.5%	4.4%	4.2%	5.6%	4.4%	14	בנים
3.3%	4.3%	3.7%	4.2%	3.6%	4.2%	4.9%	4.7%	2.9%	1.2%	6.1%	6.4%	3.4%	3.7%	4.7%	4.2%	15	בנים
2.1%	2.5%	3.2%	2.9%	2.7%	3.1%	2.9%	3.2%	4.0%	3.3%	3.4%	4.0%	2.7%	3.1%	2.9%	2.9%	14	בנות
1.7%	2.2%	2.6%	2.0%	1.8%	2.5%	3.1%	3.1%	NA*	4.6%	3.2%	2.4%	2.2%	2.4%	3.0%	2.6%	15	בנות
שיעור ההמצאות של עודף משקל בקרב בני 15-14																	
19.9%	19.6%	19.2%	20.6%	16.9%	17.7%	17.9%	18.2%	17.5%	17.9%	18.7%	16.7%	15.9%	18.0%	18.9%	19.9%	14	בנים
19.1%	19.8%	18.8%	20.0%	18.0%	18.2%	18.1%	18.8%	31.4%	13.8%	21.6%	17.1%	18.0%	18.0%	18.2%	18.1%	15	בנים
19.8%	19.4%	18.9%	21.1%	18.0%	19.8%	19.2%	19.5%	20.5%	19.8%	19.2%	19.0%	18.1%	18.6%	19.3%	18.9%	14	בנות
17.7%	17.3%	19.9%	20.8%	16.8%	18.8%	18.5%	19.0%	NA*	21.8%	20.6%	14.3%	16.2%	18.0%	18.9%	19.2%	15	בנות
שיעור ההמצאות של השמנת יתר בקרב בני 15-14																	
8.8%	15.1%	14.4%	16.5%	9.4%	14.6%	17.5%	20.9%	12.9%	15.7%	16.7%	16.5%	9.8%	13.3%	17.0%	16.6%	14	בנים
8.6%	12.9%	13.6%	15.5%	8.8%	12.4%	15.6%	17.2%	5.7%	12.5%	12.8%	12.9%	9.1%	13.5%	16.1%	18.6%	15	בנים
10.3%	12.3%	11.7%	12.3%	9.9%	14.2%	15.5%	15.0%	11.1%	14.6%	14.1%	12.9%	8.5%	12.5%	13.8%	13.7%	14	בנות
7.3%	11.2%	9.8%	10.9%	8.2%	12.1%	13.1%	12.7%	NA*	13.8%	9.5%	11.1%	8.0%	12.7%	13.8%	13.8%	15	בנות

תא מסומן ב-* מייצג ערך מכנה נמוך מ 30 מבטוחים

מבוגרים בני 65 ומעלה																
מאוחדת				מכבי				לאומית				כללית				← SES ↓ קבוצת גיל
8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	
שיעור קבלת חיסון נגד שפעת בקרב בני 65 ומעלה																
46.7%	39.5%	34.3%	35.9%	46.1%	36.8%	30.8%	29.4%	64.4%	63.3%	62.5%	59.2%	54.5%	48.2%	45.4%	49.2%	65-69
60.1%	51.4%	44.4%	41.7%	57.9%	46.8%	39.6%	37.2%	74.8%	71.6%	66.7%	68.7%	67.6%	61.0%	55.5%	57.7%	70-74
69.2%	58.1%	49.1%	47.5%	64.9%	53.2%	45.1%	41.8%	82.4%	75.1%	71.9%	69.5%	74.6%	68.9%	62.0%	61.3%	75-79
70.4%	63.9%	57.1%	53.8%	63.9%	54.3%	44.5%	41.2%	77.7%	73.9%	71.4%	70.8%	78.9%	70.9%	65.4%	64.8%	80-84
72.8%	63.4%	56.3%	56.2%	61.6%	53.8%	44.9%	39.2%	78.2%	73.5%	67.9%	68.2%	79.0%	72.9%	67.5%	65.0%	85+
45.3%	35.9%	31.1%	32.0%	45.1%	34.7%	27.6%	27.0%	69.4%	64.8%	64.2%	64.6%	51.9%	43.9%	40.4%	46.1%	65-69
55.7%	46.6%	38.4%	37.8%	55.1%	42.8%	35.2%	32.2%	75.5%	71.3%	68.6%	68.6%	63.4%	54.7%	48.7%	51.0%	70-74
63.1%	52.1%	43.7%	40.7%	59.1%	46.6%	38.6%	35.3%	78.5%	73.1%	68.8%	69.7%	69.4%	62.0%	54.5%	54.8%	75-79
65.8%	54.2%	46.2%	47.3%	56.1%	46.1%	36.9%	33.9%	74.7%	71.1%	69.1%	70.1%	70.8%	62.2%	58.2%	58.6%	80-84
63.5%	57.6%	49.2%	50.3%	49.4%	41.1%	36.5%	34.9%	69.6%	67.8%	66.9%	66.7%	71.8%	65.5%	61.5%	60.7%	85+
שיעור התייעוד של מרכיבי BMI בקרב בני 65 ומעלה																
68.0%	68.7%	69.0%	71.1%	55.0%	59.1%	60.4%	62.6%	69.1%	67.6%	68.1%	64.7%	70.4%	73.1%	73.5%	74.3%	65-69
75.1%	75.1%	72.5%	76.0%	61.3%	66.5%	65.8%	68.2%	76.1%	76.1%	72.8%	73.9%	81.4%	82.1%	80.9%	83.1%	70-74
78.2%	76.0%	74.3%	76.9%	65.1%	69.9%	68.6%	70.5%	57.9%	58.6%	56.3%	58.8%	83.7%	84.7%	82.9%	82.5%	75-79
49.3%	48.6%	45.8%	50.4%	65.3%	68.7%	66.4%	70.5%	57.2%	53.3%	51.8%	53.1%	84.8%	84.9%	83.0%	81.7%	80-84
70.1%	69.3%	70.1%	75.5%	53.5%	60.3%	61.3%	64.7%	70.7%	68.3%	70.7%	71.2%	71.1%	73.3%	75.2%	78.2%	65-69
75.6%	74.0%	73.2%	77.2%	59.8%	64.5%	65.5%	67.5%	76.9%	77.2%	75.6%	75.3%	80.6%	82.0%	81.6%	83.6%	70-74
76.1%	74.4%	71.7%	73.2%	61.8%	66.3%	65.9%	68.1%	55.9%	55.3%	54.2%	56.7%	82.7%	84.0%	82.6%	82.5%	75-79
46.8%	42.4%	44.7%	44.4%	59.9%	63.0%	63.1%	66.9%	48.3%	50.5%	49.1%	50.1%	81.6%	81.8%	81.0%	79.7%	80-84
שיעור המבוגרים בני 65 ומעלה שחלה ירידה חדה במשקלם בשנתיים האחרונות																
5.8%	6.5%	6.9%	4.1%	6.6%	7.6%	6.7%	8.4%	5.5%	6.4%	5.9%	7.7%	4.8%	5.4%	5.6%	5.1%	65-69
6.3%	6.2%	7.1%	7.8%	6.2%	7.3%	7.7%	7.3%	6.2%	6.7%	6.4%	7.4%	4.9%	5.1%	5.8%	5.2%	70-74
7.5%	7.7%	7.8%	9.3%	7.2%	9.0%	8.7%	8.8%	8.2%	7.1%	10.6%	9.9%	5.3%	5.9%	6.4%	6.9%	75-79
8.1%	8.1%	9.5%	9.5%	9.8%	9.1%	10.6%	10.3%	10.2%	11.0%	9.2%	10.2%	6.3%	7.1%	7.1%	7.5%	80-84
10.5%	11.0%	7.2%	14.5%	12.6%	12.4%	13.4%	16.2%	11.0%	8.5%	8.1%	12.1%	8.4%	8.5%	9.1%	8.5%	85+
7.3%	7.8%	8.4%	6.2%	9.1%	9.3%	8.9%	8.3%	7.9%	7.3%	7.7%	7.2%	6.5%	6.7%	6.8%	6.2%	65-69
6.5%	8.4%	8.0%	7.9%	8.6%	9.4%	8.7%	9.0%	5.9%	7.6%	7.9%	8.9%	6.2%	6.8%	6.5%	6.4%	70-74
8.9%	8.0%	10.1%	8.9%	8.9%	9.4%	10.0%	10.4%	9.8%	8.2%	9.2%	9.6%	6.3%	6.9%	7.4%	7.7%	75-79
8.6%	9.1%	10.8%	9.6%	11.6%	10.5%	11.9%	12.8%	9.8%	10.1%	10.5%	9.9%	7.4%	8.0%	8.6%	8.5%	80-84
8.2%	13.4%	12.8%	9.0%	15.4%	14.3%	15.6%	15.2%	7.9%	10.4%	14.8%	13.5%	8.8%	9.4%	9.4%	9.6%	85+
שיעור תת משקל בקרב מבוגרים בני 75 ומעלה																
11.9%	11.3%	12.3%	13.0%	11.3%	11.1%	11.0%	10.8%	13.2%	12.1%	12.5%	16.2%	13.1%	13.5%	14.1%	15.9%	75-79
17.3%	13.2%	13.4%	17.0%	15.1%	13.4%	14.4%	15.7%	16.8%	15.5%	17.8%	15.7%	17.2%	16.8%	17.1%	17.9%	80-84
18.8%	18.2%	14.9%	19.6%	20.3%	18.3%	19.7%	17.0%	20.8%	21.2%	18.3%	20.5%	20.7%	21.0%	20.9%	22.0%	85+
22.6%	17.3%	12.4%	8.7%	21.1%	15.4%	12.0%	10.3%	16.5%	14.0%	12.6%	13.2%	20.5%	16.8%	13.9%	11.0%	75-79
23.9%	18.8%	14.6%	11.5%	23.3%	16.6%	13.6%	13.0%	21.7%	18.7%	14.9%	13.3%	22.1%	18.6%	15.7%	12.3%	80-84
26.5%	21.9%	18.5%	19.6%	26.8%	19.8%	17.2%	19.5%	22.2%	19.0%	15.8%	20.5%	23.6%	21.8%	20.4%	17.6%	85+
שיעור ההמצאות של השנתון יותר בקרב בני 65 ומעלה																
26.3%	31.1%	32.2%	35.7%	25.2%	30.8%	33.5%	37.0%	26.6%	28.9%	32.6%	34.3%	25.6%	28.5%	29.4%	31.0%	65-69
23.6%	29.1%	32.6%	35.5%	23.9%	28.1%	32.3%	35.6%	25.4%	29.3%	30.3%	33.4%	24.2%	26.5%	27.7%	29.3%	70-74
22.1%	25.4%	30.4%	31.7%	21.5%	26.2%	29.2%	30.4%	23.6%	26.5%	29.4%	27.5%	21.1%	24.0%	26.1%	28.1%	75-79
18.8%	23.5%	24.3%	29.4%	17.6%	21.8%	23.7%	25.0%	19.4%	20.6%	23.2%	27.1%	16.9%	20.2%	21.7%	24.6%	80-84
24.1%	32.6%	42.1%	53.8%	24.4%	33.9%	42.4%	50.9%	25.5%	33.4%	43.5%	49.1%	26.6%	33.4%	41.6%	53.7%	65-69
25.1%	32.9%	41.1%	52.1%	25.7%	35.5%	42.2%	48.8%	27.0%	34.1%	43.0%	49.2%	25.8%	32.8%	40.3%	50.6%	70-74
23.3%	32.7%	40.0%	48.6%	23.6%	34.4%	41.5%	47.0%	28.0%	35.2%	43.4%	48.0%	25.4%	31.7%	38.5%	47.3%	75-79
21.9%	31.4%	37.6%	43.4%	22.7%	31.8%	38.6%	42.8%	24.8%	32.1%	36.1%	39.6%	23.3%	29.7%	35.1%	43.1%	80-84
שיעור השימוש בבוזיאופנים ארוכי טווח בקרב בני 65 ומעלה																
0.9%	1.1%	1.6%	1.5%	1.0%	1.4%	1.7%	1.5%	1.5%	2.1%	2.3%	2.4%	2.2%	2.6%	2.6%	2.2%	65-69
1.5%	1.5%	2.5%	1.9%	1.5%	1.6%	1.9%	1.6%	2.9%	3.0%	3.4%	3.5%	3.4%	3.6%	3.6%	2.7%	70-74
2.3%	1.6%	2.9%	2.0%	1.9%	2.0%	2.3%	2.1%	3.7%	4.0%	4.8%	3.5%	5.2%	5.2%	4.5%	3.2%	75-79
2.8%	3.7%	3.5%	2.3%	3.0%	3.1%	3.5%	3.0%	7.6%	7.5%	8.4%	5.4%	7.8%	7.7%	7.0%	4.7%	80-84
4.0%	4.3%	6.4%	4.8%	3.3%	4.1%	4.1%	4.1%	8.3%	8.8%	10.4%	9.3%	11.4%	10.7%	9.4%	6.9%	85+
1.1%	1.7%	2.5%	1.7%	1.9%	2.0%	2.7%	2.4%	2.8%	3.4%	3.6%	2.7%	3.3%	3.4%	3.2%	2.2%	65-69
2.0%	2.3%	3.0%	2.1%	2.6%	2.8%	2.9%	3.1%	4.4%	4.2%	5.1%	3.4%	5.0%	4.7%	4.4%	3.1%	70-74
3.3%	3.2%	3.5%	3.1%	3.1%	3.5%	3.5%	3.5%	5.6%	6.3%	5.8%	5.7%	7.3%	6.9%	6.2%	3.4%	75-79
4.1%	4.6%	5.2%	4.8%	4.4%	4.6%	5.0%	5.0%	8.7%	10.7%	8.4%	9.9%	11.2%	10.6%	8.8%	6.3%	80-84
5.9%	6.6%	8.4%	6.3%	5.4%	5.4%	5.9%	4.9%	12.8%	12.5%	13.2%	15.2%	16.6%	14.7%	13.5%	8.9%	85+
שיעור השימוש בבוזיאופנים ארוכי טווח בקרב בני 65 ומעלה																
1.1%	1.3%	1.0%	1.8%	1.1%	1.2%	1.2%	1.8%	1.4%	0.9%	1.4%	1.9%	1.1%	1.2%	1.5%	1.7%	65-69
1.5%	1.3%	1.4%	2.0%	1.4%	1.1%	1.3%	1.4%	1.4%	1.0%	1.4%	1.7%	1.2%	1.2%	1.4%	1.3%	70-74
2.0%	1.7%	1.5%	1.6%	1.7%	1.4%	1.1%	1.8%	0.8%	1.4%	1.3%	1.7%	1.4%	1.3%	1.4%	1.3%	75-79
2.1%	1.7%	1.4%	1.5%	1.9%	1.3%	1.3%	1.4%	1.8%	1.5%	1.4%	2.1%	1.3%	1.5%	1.3%	1.2%	80-84
2.0%	1.5%	1.4%	2.5%	1.5%	1.1%	1.1%	1.0%	1.3%	1.2%	1.6%	1.9%	1.8%	1.3%	1.1%	1.2%	85+
1.6%	1.6%	1.7%	2.5%	1.9%	1.7%	1.5%	1.7%	1.3%	1.4%	1.4%	1.0%	1.6%	1.5%	1.3%	1.4%	65-69
1.9%	1.8%	1.7%	1.8%	2.2%	1.8%	1.6%	1.5%	2.0%	1.4%	1.5%	1.6%	2.0%	1.6%	1.5%	1.2%	70-74
2.4%	2.3%	2.2%	2.2%	2.8%	2.1%	1.6%	1.9%	2.5%	1.3%	1.9%	1.9%	2.1%	1.9%	1.7%	1.4%	75-79
2.6%	2.0%	2.0%	2.4%	2.2%	2.0%	1.6%	2.1%	1.9%	1.7%	1.6%	1.8%	2.4%	2.1%	1.7%	1.5%	80-84
2.5%	1.7%	1.8%	1.3%	2.4%	1.6%	1.6%	1.7%	2.3%	1.6%	1.9%	1.6%	2.0%	2.0%	1.8%	1.3%	85+

חא מסומן ב- * מייצג ערך מכנה נמוך מ-30 מבוגרים

מחלות נשימתיות																
מאוחדת				מבני				לאומית				כללית				← SES ↓ קבוצת גיל ↓ מין
8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	
שיעור ההמצאות של אסתמה ממשכת בקרב בני 44-5																
1.3%	1.0%	0.7%	0.6%	1.2%	1.2%	1.0%	0.7%	1.9%	1.7%	1.0%	0.6%	0.9%	1.0%	0.9%	0.8%	5-14
1.0%	0.8%	0.5%	0.4%	0.9%	0.8%	0.5%	0.4%	1.3%	1.1%	0.6%	0.5%	0.8%	0.6%	0.5%	0.4%	15-24
1.2%	1.1%	0.8%	0.6%	0.9%	0.8%	0.7%	0.6%	1.5%	1.3%	0.9%	0.6%	0.7%	0.7%	0.6%	0.6%	25-34
1.6%	1.3%	1.0%	1.0%	1.1%	1.0%	1.0%	0.8%	1.6%	1.5%	1.2%	1.1%	1.0%	1.0%	0.9%	1.0%	35-44
0.8%	0.7%	0.4%	0.3%	0.8%	0.8%	0.6%	0.5%	0.9%	0.9%	0.6%	0.3%	0.6%	0.6%	0.5%	0.5%	5-14
0.9%	0.7%	0.4%	0.3%	0.7%	0.6%	0.5%	0.3%	1.3%	0.9%	0.5%	0.3%	0.6%	0.5%	0.4%	0.3%	15-24
1.1%	1.0%	0.6%	0.4%	0.8%	0.9%	0.8%	0.5%	1.5%	1.2%	0.9%	0.8%	0.7%	0.8%	0.6%	0.6%	25-34
1.3%	1.5%	1.0%	0.9%	1.0%	1.2%	1.2%	0.9%	1.4%	1.8%	1.3%	1.3%	1.0%	1.0%	1.1%	1.0%	35-44
שיעור יחס התחפות לאסתמה (AMR) גדול או שווה 0.5 בקרב בני 44-5																
82.4%	82.9%	85.2%	89.7%	75.8%	72.7%	72.2%	83.3%	76.5%	75.0%	75.4%	78.1%	77.3%	77.6%	75.1%	80.5%	5-14
80.5%	75.0%	81.4%	81.3%	79.8%	78.5%	68.1%	82.4%	73.7%	65.9%	75.5%	71.1%	85.5%	77.8%	74.8%	81.2%	15-24
74.2%	70.8%	80.4%	85.6%	77.4%	74.5%	75.6%	75.5%	73.2%	66.1%	74.3%	71.6%	80.6%	80.0%	77.8%	76.2%	25-34
79.3%	75.9%	77.0%	86.7%	72.8%	72.4%	67.2%	81.6%	71.0%	64.8%	75.0%	79.0%	75.9%	76.6%	74.7%	76.1%	35-44
84.3%	81.3%	82.1%	88.9%	72.8%	72.0%	78.3%	86.0%	73.7%	73.3%	79.5%	77.8%	76.1%	75.8%	73.1%	81.4%	5-14
79.1%	71.8%	82.4%	88.8%	79.9%	77.7%	72.8%	76.9%	66.7%	65.8%	73.3%	69.6%	84.2%	82.2%	79.1%	83.5%	15-24
81.4%	78.0%	84.2%	87.5%	80.0%	79.2%	72.9%	76.0%	78.8%	64.6%	78.0%	81.8%	80.3%	79.6%	76.9%	80.1%	25-34
76.4%	78.8%	84.0%	88.1%	79.4%	75.0%	70.8%	81.4%	78.4%	76.5%	78.6%	79.3%	79.6%	78.2%	77.1%	81.0%	35-44
שיעור קבלת חיסון נגד שפעת בקרב חולי אסתמה ממשכת בני 44-5																
20.8%	20.1%	14.8%	16.7%	30.8%	21.0%	16.9%	15.2%	29.6%	38.5%	44.0%	41.1%	32.0%	23.7%	22.7%	33.2%	5-14
18.2%	15.3%	14.7%	12.3%	19.6%	15.3%	8.8%	13.9%	31.6%	45.1%	33.0%	38.6%	20.5%	19.1%	18.4%	23.5%	15-24
11.3%	14.6%	9.5%	15.0%	17.3%	13.0%	10.2%	10.0%	30.4%	33.9%	27.9%	37.5%	19.4%	13.6%	17.4%	27.4%	25-34
20.7%	18.1%	17.3%	14.9%	20.7%	18.3%	15.7%	13.6%	35.5%	28.1%	44.1%	41.9%	23.3%	18.4%	27.2%	36.7%	35-44
27.0%	18.1%	19.6%	25.3%	28.3%	24.8%	19.0%	16.9%	39.5%	45.5%	47.0%	43.1%	30.0%	27.3%	21.8%	33.5%	5-14
13.4%	14.5%	9.9%	17.3%	18.8%	16.7%	19.6%	12.8%	41.7%	52.6%	44.2%	44.9%	21.0%	22.6%	19.2%	33.8%	15-24
23.3%	21.3%	15.8%	21.2%	24.5%	18.4%	14.6%	15.0%	53.8%	55.6%	57.6%	59.6%	26.4%	19.7%	20.4%	31.0%	25-34
22.7%	19.1%	18.2%	13.1%	23.5%	17.0%	17.9%	14.2%	41.2%	51.7%	51.6%	58.6%	27.8%	21.8%	23.0%	34.3%	35-44
שיעור הביצוע של ספירומטריה בקרב חולי COPD ואנולוסייה בסיכון גבוה ל-COPD בני 74-50																
79.5%	77.8%	75.7%	72.4%	55.4%	59.6%	63.2%	61.2%	71.8%	66.4%	75.0%	69.6%	69.4%	72.1%	70.9%	76.5%	50-54
74.3%	74.1%	80.0%	76.7%	62.2%	67.1%	64.1%	58.9%	77.5%	70.7%	76.9%	73.9%	79.8%	78.3%	74.9%	76.4%	55-59
78.5%	76.0%	81.9%	81.5%	72.6%	71.5%	69.5%	63.2%	82.7%	76.5%	80.5%	80.6%	77.3%	80.1%	80.9%	79.7%	60-64
84.5%	83.2%	83.8%	78.6%	70.5%	73.7%	70.8%	63.7%	84.9%	83.0%	85.5%	80.2%	82.9%	84.9%	80.6%	78.5%	65-69
76.7%	87.4%	81.5%	82.8%	67.6%	77.1%	72.8%	64.8%	85.6%	85.3%	85.7%	80.0%	87.9%	85.0%	84.0%	79.5%	70-74
62.5%	74.2%	74.2%	69.1%	54.8%	58.5%	58.5%	51.7%	67.7%	65.0%	74.4%	68.3%	69.8%	71.3%	73.2%	69.3%	50-54
81.7%	75.4%	76.2%	73.5%	63.1%	66.6%	62.7%	54.3%	69.8%	70.8%	74.6%	78.9%	74.5%	78.4%	77.6%	76.3%	55-59
78.4%	83.3%	81.5%	75.5%	66.4%	68.8%	66.7%	54.1%	73.3%	78.6%	77.1%	79.5%	82.2%	82.5%	82.0%	77.1%	60-64
77.3%	81.4%	81.8%	75.4%	71.1%	69.5%	70.2%	64.9%	81.1%	78.9%	82.8%	79.1%	83.7%	82.9%	81.5%	79.2%	65-69
83.9%	83.0%	85.1%	69.6%	70.6%	75.2%	71.3%	65.6%	79.5%	86.4%	80.0%	84.3%	87.2%	83.4%	82.6%	76.6%	70-74
חיסון ערכי בדיקת ספירומטריה בקרב חולי COPD ואנולוסייה בסיכון גבוה ל-COPD בני 74-50																
0.0%	0.0%	0.0%	0.0%	38.2%	31.9%	28.6%	13.9%	60.8%	48.6%	43.7%	42.3%	60.5%	56.5%	57.7%	62.5%	50-54
0.0%	0.0%	0.0%	0.0%	36.9%	35.1%	30.4%	25.3%	60.0%	55.0%	52.6%	40.5%	51.6%	66.5%	64.2%	65.3%	55-59
0.0%	0.0%	0.0%	0.0%	41.6%	36.4%	29.5%	16.7%	56.7%	52.7%	58.0%	48.0%	59.9%	68.5%	63.7%	67.1%	60-64
0.0%	0.0%	0.0%	0.0%	42.3%	32.7%	32.2%	22.3%	61.2%	53.3%	53.1%	42.4%	60.9%	67.9%	68.1%	69.1%	65-69
0.0%	0.0%	0.0%	0.0%	43.5%	34.7%	34.4%	13.1%	59.0%	51.0%	51.4%	54.1%	58.7%	66.7%	65.8%	68.9%	70-74
0.0%	0.0%	0.0%	0.0%	42.4%	38.0%	34.4%	17.7%	42.9%	49.6%	44.9%	31.4%	55.2%	58.6%	64.5%	67.9%	50-54
0.0%	0.0%	0.0%	0.0%	44.5%	38.1%	30.8%	17.1%	56.7%	49.5%	56.2%	41.3%	62.2%	64.0%	69.2%	62.3%	55-59
0.0%	0.0%	0.0%	0.0%	45.8%	33.3%	35.2%	19.8%	56.8%	49.3%	50.0%	44.7%	59.9%	63.9%	64.8%	62.5%	60-64
0.0%	0.0%	0.0%	0.0%	47.9%	39.7%	33.6%	17.7%	64.2%	48.6%	51.5%	46.4%	57.8%	60.7%	66.8%	62.3%	65-69
0.0%	0.0%	0.0%	0.0%	48.2%	44.9%	37.2%	17.8%	53.5%	48.9%	50.0%	45.1%	59.1%	64.0%	66.3%	65.6%	70-74

חא מסומן ב- * מייצג ערך מכנה נמוך מ-30 מבטוחים

לב וכולי דם																
מאוחדת				מכבי				לאומית				כללית				← SES ↓ מיץ ↓ קבוצת גיל
8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	
שיעור הביצוע של בדיקת כולסטרול בקרב בני 54-35																
78.0%	76.5%	73.4%	71.5%	80.0%	78.0%	74.9%	73.1%	78.9%	74.4%	73.7%	70.4%	82.6%	79.1%	75.8%	74.3%	35-39
80.2%	79.5%	78.1%	77.9%	82.2%	81.3%	79.0%	79.4%	79.3%	78.3%	78.3%	76.2%	84.5%	82.2%	79.8%	79.8%	40-44
83.9%	83.3%	81.9%	82.3%	85.9%	84.9%	83.5%	83.8%	83.7%	82.8%	82.3%	82.3%	87.1%	85.6%	84.1%	84.1%	45-49
88.5%	88.1%	86.9%	87.4%	90.2%	89.5%	87.8%	88.1%	89.6%	87.1%	85.7%	86.8%	91.1%	90.0%	89.1%	88.8%	50-54
88.4%	88.0%	85.8%	86.3%	86.5%	86.7%	85.5%	86.0%	88.7%	88.2%	87.3%	83.5%	92.0%	91.1%	90.7%	91.2%	35-39
90.6%	89.8%	89.4%	89.4%	88.8%	88.7%	87.9%	88.1%	90.7%	90.4%	89.8%	87.7%	92.5%	92.1%	92.0%	92.8%	40-44
92.7%	92.6%	91.7%	92.5%	91.3%	91.7%	91.1%	90.6%	92.9%	91.9%	92.2%	90.7%	93.7%	93.6%	93.3%	94.3%	45-49
94.5%	94.4%	93.5%	94.3%	94.0%	94.1%	93.3%	94.0%	94.6%	94.1%	94.7%	93.7%	95.3%	95.3%	95.1%	96.0%	50-54
שיעור הביצוע של בדיקת כולסטרול בקרב בני 74-55																
61.3%	61.1%	59.6%	64.5%	55.6%	57.7%	56.8%	59.2%	63.0%	62.1%	62.1%	60.0%	64.8%	64.6%	64.4%	65.8%	55-59
69.4%	68.7%	68.3%	70.6%	63.7%	65.3%	64.8%	65.1%	72.6%	69.8%	67.8%	67.6%	73.0%	72.5%	70.5%	71.8%	60-64
75.8%	75.6%	74.2%	77.0%	71.5%	73.2%	72.6%	71.8%	78.8%	78.8%	75.4%	72.5%	80.4%	80.0%	77.7%	78.1%	65-69
81.6%	81.2%	79.5%	81.4%	77.7%	79.7%	77.7%	76.5%	82.1%	82.3%	80.1%	78.7%	85.3%	84.5%	82.5%	82.7%	70-74
69.1%	68.7%	69.1%	74.4%	54.9%	57.8%	57.9%	62.4%	70.3%	71.3%	71.8%	71.6%	69.3%	70.8%	72.5%	76.8%	55-59
72.8%	72.8%	72.8%	77.4%	62.1%	64.9%	66.4%	69.0%	75.6%	77.8%	76.6%	76.0%	75.2%	76.3%	77.5%	80.6%	60-64
79.1%	77.8%	78.3%	81.8%	69.2%	72.0%	73.0%	73.0%	80.3%	82.0%	80.5%	80.9%	81.5%	82.0%	81.8%	84.5%	65-69
80.9%	81.9%	81.6%	84.6%	74.6%	77.5%	77.1%	77.6%	85.0%	84.2%	83.9%	81.8%	85.0%	85.8%	85.1%	86.0%	70-74
שיעור עמידה ביעד LDL לפי קבוצת סיכון למחלות לב בקרב בני 74-35																
87.0%	83.0%	81.9%	80.5%	88.2%	85.3%	84.8%	86.1%	84.0%	81.5%	81.4%	78.6%	92.4%	91.0%	90.0%	88.8%	35-44
77.1%	71.4%	71.2%	69.5%	80.6%	77.3%	75.1%	78.4%	72.2%	67.8%	65.6%	66.4%	83.3%	79.7%	78.1%	76.3%	45-54
77.4%	74.0%	73.8%	72.7%	84.4%	80.3%	78.1%	79.4%	74.3%	69.4%	68.6%	68.3%	86.7%	83.7%	81.7%	79.8%	55-64
80.6%	77.5%	74.7%	76.0%	87.2%	82.3%	80.6%	80.9%	78.0%	73.0%	71.4%	70.2%	90.9%	88.2%	86.6%	85.4%	65-74
94.4%	93.6%	94.0%	94.3%	95.3%	93.3%	93.6%	94.2%	92.1%	91.3%	92.5%	93.3%	96.7%	96.5%	96.5%	96.3%	35-44
88.9%	87.7%	87.8%	89.0%	90.3%	88.2%	87.7%	89.2%	86.6%	84.4%	85.5%	87.1%	92.5%	91.0%	91.2%	91.5%	45-54
79.6%	76.2%	76.5%	79.9%	84.2%	80.7%	78.7%	82.7%	76.0%	72.6%	72.9%	74.3%	86.5%	83.9%	83.9%	85.5%	55-64
82.2%	82.1%	79.1%	82.6%	86.9%	84.8%	82.6%	84.0%	79.8%	76.6%	76.5%	77.3%	90.5%	89.1%	88.8%	89.4%	65-74
שיעור בעלי תמת LDL נמוכה או שווה ל-160 מ"ג/ל לפי קבוצת סיכון נמוך למחלות לב וכולי דם בבני 74-35																
91.5%	89.2%	89.1%	89.8%	91.4%	89.5%	88.7%	90.7%	88.1%	85.9%	86.8%	86.4%	93.1%	92.2%	91.9%	91.0%	35-44
88.8%	87.5%	87.0%	88.8%	89.5%	87.8%	86.9%	88.8%	84.1%	84.2%	82.3%	85.3%	90.2%	88.9%	89.2%	88.8%	45-54
91.2%	89.4%	89.9%	89.5%	91.9%	90.5%	88.9%	90.7%	85.3%	84.5%	83.9%	87.2%	92.4%	91.3%	90.8%	91.3%	55-64
94.1%	91.4%	92.6%	93.5%	94.6%	92.1%	91.0%	91.9%	87.5%	88.5%	90.0%	87.3%	95.5%	94.5%	94.3%	94.6%	65-74
95.0%	94.5%	94.8%	95.1%	95.7%	94.0%	94.5%	94.8%	92.6%	92.6%	93.2%	94.2%	96.8%	96.8%	96.8%	96.5%	35-44
90.6%	89.5%	89.7%	90.9%	91.3%	90.0%	90.2%	91.1%	87.7%	86.0%	87.1%	88.5%	92.9%	91.8%	92.1%	92.9%	45-54
86.8%	84.9%	85.5%	87.5%	89.3%	87.9%	87.1%	89.1%	83.9%	81.8%	82.7%	84.6%	90.3%	89.7%	89.9%	90.5%	55-64
88.1%	89.0%	87.7%	91.0%	91.4%	90.5%	88.8%	89.6%	84.9%	85.8%	84.2%	85.2%	93.1%	92.2%	92.4%	93.9%	65-74
שיעור בעלי תמת LDL נמוכה או שווה ל-130 מ"ג/ל בקרב בני 2 או יותר נקודות גרמי סיכון למחלות לב וכולי דם - Framingham נמוך או שווה ל-20% בבני 74-35																
61.8%	57.9%	60.6%	60.7%	68.0%	66.6%	70.1%	71.4%	62.4%	60.8%	63.6%	58.9%	80.9%	79.5%	77.6%	78.5%	35-44
64.2%	59.7%	63.7%	62.8%	70.7%	70.4%	70.0%	75.2%	62.6%	59.5%	60.3%	60.6%	74.1%	72.5%	72.6%	72.6%	45-54
68.1%	67.0%	69.4%	70.3%	79.7%	78.0%	78.3%	78.4%	71.1%	67.5%	69.8%	69.5%	82.9%	82.2%	81.2%	81.0%	55-64
79.0%	76.8%	74.5%	75.6%	88.1%	84.1%	83.4%	82.5%	81.4%	74.9%	74.6%	74.8%	90.0%	87.7%	86.0%	85.3%	65-74
61.8%	65.4%	68.2%	64.0%	73.4%	71.5%	70.2%	70.3%	78.9%	62.7%	73.3%	67.8%	85.1%	84.1%	85.6%	85.9%	35-44
50.7%	57.9%	61.9%	62.5%	65.5%	62.2%	68.9%	62.9%	62.9%	58.2%	60.9%	64.0%	76.0%	74.6%	75.1%	76.6%	45-54
56.6%	55.9%	58.9%	66.4%	65.2%	63.4%	63.2%	71.1%	56.8%	55.7%	57.2%	59.7%	73.4%	70.8%	72.8%	77.7%	55-64
69.4%	70.4%	67.5%	72.1%	76.8%	76.2%	74.4%	77.0%	71.9%	65.4%	68.5%	69.9%	83.2%	82.8%	82.9%	83.3%	65-74
שיעור היתעד של מדידת לחץ דם בקרב בני 54-20																
70.8%	67.0%	63.9%	68.8%	74.5%	74.2%	66.8%	66.1%	58.2%	62.4%	65.7%	65.9%	75.4%	74.0%	72.4%	76.0%	20-24
70.9%	70.9%	68.8%	73.0%	77.0%	79.1%	76.4%	72.7%	69.1%	71.7%	71.7%	71.9%	74.5%	76.1%	75.9%	78.5%	25-34
81.4%	80.0%	78.3%	80.7%	83.9%	84.5%	82.2%	80.4%	73.8%	75.4%	76.0%	76.9%	80.5%	81.8%	82.4%	85.5%	35-44
91.2%	90.2%	89.3%	90.7%	89.7%	90.4%	89.3%	89.1%	81.9%	83.5%	83.8%	86.5%	90.4%	91.8%	92.8%	94.8%	45-54
83.9%	81.1%	83.1%	86.5%	83.0%	85.0%	83.8%	84.1%	75.4%	80.1%	85.8%	85.5%	84.8%	84.9%	84.2%	86.9%	20-24
87.5%	90.0%	92.4%	95.0%	89.8%	92.1%	92.0%	92.4%	87.2%	89.7%	90.9%	92.4%	88.4%	90.6%	90.4%	90.9%	25-34
91.1%	90.2%	92.9%	94.5%	90.5%	91.2%	90.9%	91.3%	84.5%	86.8%	88.6%	91.5%	89.2%	90.4%	91.2%	92.8%	35-44
93.8%	93.4%	93.1%	95.5%	90.7%	92.1%	92.3%	92.5%	85.7%	87.5%	89.0%	89.9%	92.7%	94.7%	95.7%	97.2%	45-54
שיעור היתעד של מדידת לחץ דם בקרב בני 74-55																
56.7%	57.5%	60.0%	65.1%	57.8%	61.8%	62.5%	65.0%	54.5%	56.5%	59.0%	60.0%	59.9%	64.4%	67.1%	71.1%	55-59
62.0%	64.4%	66.0%	71.4%	65.9%	69.2%	68.6%	71.8%	60.4%	63.0%	62.9%	67.6%	69.0%	72.6%	72.9%	76.4%	60-64
73.4%	72.9%	73.1%	76.4%	74.0%	76.6%	76.4%	77.5%	70.6%	72.7%	70.5%	70.5%	77.0%	79.9%	79.9%	82.5%	65-69
79.2%	80.2%	77.9%	82.4%	80.7%	82.7%	81.4%	81.3%	74.8%	76.8%	75.1%	76.9%	83.4%	85.0%	84.1%	86.1%	70-74
57.8%	58.4%	63.0%	70.6%	55.5%	60.5%	62.6%	67.8%	56.1%	62.6%	65.1%	66.6%	60.5%	65.9%	71.3%	77.0%	55-59
63.4%	64.9%	67.9%	75.0%	63.4%	68.3%	70.3%	72.5%	64.4%	66.9%	69.0%	69.8%	68.1%	73.1%	75.7%	81.3%	60-64
73.8%	73.7%	74.5%	80.1%	72.3%	76.8%	79.9%	71.3%	71.3%	72.7%	73.9%	75.7%	77.1%	79.9%	81.5%	85.4%	65-69
79.0%	78.5%	77.9%	83.7%	78.5%	81.6%	82.1%	83.1%	76.2%	76.6%	76.7%	76.1%	82.3%	84.6%	84.8%	87.5%	70-74
שיעור בעלי תמת LDL נמוכה או שווה ל-70 מ"ג/ל או טיפול בטסטטים בעמידת גבוהה לאחר צנתור לבני / ניתוח מעקפים בקרב בני 79-35																
80.6%	86.2%	78.9%	83.5%	79.0%	76.0%	68.3%	68.2%	60.0%	64.1%	61.2%	49.4%	76.5%	73.4%	66.8%	62.3%	35-49
84.6%	85.8%	84.1%	84.4%	79.8%	78.3%	69.7%	68.3%	77.6%	68.8%	63.5%	65.6%	78.9%	75.9%	70.0%	66.6%	50-59
89.5%	85.8%	84.7%	85.1%	82.0%	77.2%	73.9%	76.7%	78.2%	74.6%	71.2%	66.2%	83.3%	79.3%	75.8%	73.0%	60-69
89.0%	88.4%	86.1%	86.6%	83.9%	81.4%	76.5%	79.0%	82.7%	79.6%	75.3%	71.3%	86.4%	83.9%	80.0%	77.9%	70-79
NA*	46.2%	37.5%	39.3%	72.7%	65.6%	72.9%	NA*	NA*	NA*	NA*	NA*	64.4%	50.9%	61.7%	54.9%	35-49
64.8%	63.0%	67.0%	73.6%	75.1%	71.4%	65.1%	63.8%	NA*	61.2%	56.1%	60.0%	66.9%	69.9%	62.6%	64.7%	50-59
78.5%	74.2%	77.0%	77.8%	76.5%	70.1%	68.4%	72.3%	68.9%	68.9%	68.6%	66.7%	76.8%	71.9%	71.8%	71.4%	60-69
81.2%	74.7%	76.8%	80.4%	77.1%	71.4%	72.4%	74.0%	74.8%	70.8%	62.9%	70.1%	77.0%	76.9%	74.9%	75.4%	70-79
שיעור הטיפול בתחנות להורדת LDL לאחר ניתוח מעקפים /או צנתור לבני בקרב בני 79-35																
69.8%	75.9%	71.2%	68.3%	79.4%	81.6%	79.1%	73.6%	86.5%	78.5%	83.3%	73.1%	76.9%	80.3%	75.5%	77.2%	35-49
78.2%	77.2%	78.5%	76.6%	86.6%	85.6%	79.9%	82.6%	88.0%	79.8%	79.8%	79.3%	83.7%	82.4%	80.6%	79.1%	50-59
84.3%	81.1%	82.0%	80.8%	89.2%	85.2%	81.2%	88.6%	86.8%	85.8%	85.0%	84.6%	87.0%	85.0%	82.0%	81.7%	60-69
88.1%	87.7%	86.5%	81.8%	90.8%	87.6%	85.3%	88.6%	92.0%	91.5%	87.9%	82.9%	90.7%	88.1%	86.4%	85.7%	70-79
NA*	45.0%	28.6%	31.4%	71.9%	66.7%	70.0%	NA*	NA*	NA*	NA*	NA*	71.9%	54.3%	76.3%	71.8%	35-49
78.0%	60.9%	68.6%	69.0%	84.7%	80.6%	83.8%	84.8%	NA*	81.4%	90.2%	73.0%	77.6%	77.0%	83.0%	82.0%	50-59
76.1%	71.5%	73.4%	77.2%	88.4%	86.1%	86.6%	89.4%	100.0%	88.7%	88.6%	74.6%	85.7%	84.7%	85.8%	85.6%	60-69
80.3%	78.1%	77.2%	81.9%	90.8%	87.6%	89.2%	88.6%	86.1%	89.8%	89.1%	9					

סוכרת																← SES	↓ מינ
מאוחדת				מכבי				לאומית				כללית					
8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3		
שיעור ההימצאות של סוכרת בקרב בני 18 ומעלה																	
0.9%	0.7%	0.5%	0.4%	0.9%	1.0%	0.8%	0.7%	0.6%	0.5%	0.5%	0.4%	1.0%	0.9%	0.6%	0.5%	18-24	
0.6%	0.7%	0.7%	1.0%	0.6%	0.9%	1.0%	1.2%	0.4%	0.7%	0.7%	0.9%	0.6%	0.7%	0.9%	1.1%	25-34	
1.5%	1.8%	2.5%	4.3%	1.8%	2.5%	3.4%	4.3%	1.8%	2.2%	2.6%	3.5%	1.4%	2.2%	3.2%	4.3%	35-44	
5.0%	7.1%	9.3%	13.1%	5.7%	8.9%	10.9%	13.1%	6.7%	8.8%	9.6%	11.8%	5.3%	8.2%	10.9%	13.9%	45-54	
13.5%	16.7%	20.7%	27.2%	14.1%	18.7%	21.4%	26.4%	17.3%	20.5%	20.8%	23.5%	16.1%	20.6%	23.8%	28.1%	55-64	
26.7%	29.3%	30.8%	39.2%	25.4%	31.1%	33.2%	36.2%	29.7%	31.7%	33.1%	33.6%	29.0%	33.2%	35.3%	39.5%	65-74	
32.1%	34.1%	35.3%	40.7%	32.5%	36.3%	36.2%	37.6%	35.6%	38.1%	35.0%	37.9%	34.9%	38.2%	40.5%	43.0%	75-84	
28.9%	30.6%	28.0%	34.9%	28.2%	30.8%	29.2%	31.5%	29.2%	29.3%	29.8%	29.4%	30.3%	32.0%	32.2%	33.2%	85+	
0.8%	0.6%	0.5%	0.4%	1.1%	1.1%	0.8%	0.6%	0.8%	0.6%	0.4%	0.3%	0.8%	0.7%	0.6%	0.5%	18-24	
0.7%	0.7%	0.7%	0.8%	1.1%	1.5%	1.7%	1.3%	0.8%	0.8%	0.7%	0.8%	0.7%	0.8%	0.9%	1.1%	25-34	
1.0%	1.2%	1.6%	2.7%	2.2%	3.4%	4.1%	4.2%	1.4%	1.7%	1.9%	2.5%	1.1%	1.6%	2.2%	3.6%	35-44	
2.6%	3.9%	5.3%	9.7%	4.9%	7.6%	9.7%	10.8%	3.6%	4.7%	7.0%	9.4%	3.2%	5.4%	7.9%	12.8%	45-54	
7.8%	10.6%	14.8%	24.7%	9.6%	14.2%	17.7%	22.9%	9.2%	12.7%	17.4%	20.8%	10.1%	14.0%	19.6%	28.8%	55-64	
17.2%	20.5%	26.2%	36.6%	17.2%	24.3%	27.7%	33.3%	19.5%	23.8%	28.5%	33.4%	19.5%	24.6%	31.0%	42.1%	65-74	
23.4%	26.8%	32.1%	39.2%	22.0%	29.4%	32.1%	36.7%	26.2%	31.4%	33.7%	37.6%	25.4%	30.9%	36.1%	43.8%	75-84	
20.3%	24.8%	28.9%	29.2%	19.9%	25.9%	28.7%	30.7%	24.5%	26.1%	28.6%	25.9%	22.3%	26.3%	29.5%	31.6%	85+	

שיעור הביצוע של בדיקת HbA1c בחולי סוכרת בני 18 ומעלה																← SES	↓ מינ
מאוחדת				מכבי				לאומית				כללית					
8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3		
70.6%	81.0%	81.6%	77.0%	74.9%	75.6%	67.7%	80.9%	NA*	NA*	61.8%	71.4%	77.2%	78.0%	75.8%	78.4%	18-24	
80.9%	75.8%	82.7%	79.4%	67.4%	60.8%	69.9%	71.7%	NA*	65.0%	79.2%	71.8%	75.7%	75.3%	78.2%	77.6%	25-34	
76.1%	77.3%	80.6%	84.2%	71.1%	71.3%	75.3%	81.0%	83.3%	70.0%	72.2%	73.4%	76.7%	76.6%	78.5%	80.2%	35-44	
81.4%	78.4%	81.3%	85.1%	78.6%	81.6%	84.5%	86.5%	77.4%	76.1%	79.1%	81.9%	80.9%	82.5%	82.6%	85.5%	45-54	
84.6%	84.8%	85.4%	88.0%	86.9%	88.2%	89.1%	90.5%	83.3%	83.7%	83.6%	84.8%	86.7%	86.5%	86.8%	87.8%	55-64	
88.4%	89.2%	89.8%	91.8%	92.8%	93.7%	93.4%	94.3%	89.6%	88.9%	89.4%	88.6%	91.1%	91.5%	91.2%	91.4%	65-74	
91.6%	92.3%	91.9%	92.3%	94.7%	95.1%	94.9%	95.2%	90.4%	90.3%	89.4%	90.2%	92.2%	92.5%	92.1%	92.3%	75-84	
89.4%	88.7%	87.9%	87.8%	93.2%	92.2%	92.3%	93.1%	90.0%	86.4%	86.3%	85.6%	89.7%	89.9%	90.7%	89.3%	85+	
80.6%	83.9%	77.8%	86.2%	64.1%	56.9%	64.1%	73.7%	NA*	75.0%	69.8%	97.6%	80.4%	83.9%	87.4%	87.6%	18-24	
80.4%	78.4%	84.0%	87.7%	56.6%	52.5%	52.9%	67.1%	NA*	66.1%	79.4%	87.6%	84.9%	77.9%	81.8%	82.8%	25-34	
81.7%	80.3%	81.3%	87.5%	55.0%	57.8%	59.8%	71.7%	76.5%	80.7%	76.7%	78.3%	77.9%	79.8%	82.9%	85.8%	35-44	
84.3%	83.8%	86.0%	92.2%	62.7%	66.9%	75.9%	82.8%	90.0%	86.1%	87.2%	87.0%	83.6%	85.9%	86.9%	89.5%	45-54	
85.6%	85.2%	88.3%	92.8%	78.0%	82.8%	85.9%	91.1%	86.5%	85.1%	87.7%	90.3%	87.6%	88.4%	89.6%	92.0%	55-64	
92.0%	91.2%	91.1%	93.7%	90.5%	92.2%	93.1%	94.4%	90.9%	91.7%	90.5%	89.3%	91.9%	92.0%	91.9%	93.3%	65-74	
92.1%	90.5%	91.4%	91.6%	93.5%	94.0%	94.6%	94.9%	90.1%	90.0%	88.5%	88.1%	92.0%	92.2%	92.2%	92.7%	75-84	
88.0%	85.6%	87.3%	87.0%	89.7%	91.8%	91.4%	92.2%	83.0%	85.7%	82.3%	84.4%	87.2%	87.9%	88.9%	89.8%	85+	

שיעור בעלי תת HbA1c מאוזנת בחולי סוכרת בני 18 ומעלה																← SES	↓ מינ
מאוחדת				מכבי				לאומית				כללית					
8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3		
69.4%	64.8%	55.3%	52.3%	74.3%	73.8%	63.8%	58.9%	NA*	71.7%	57.4%	48.5%	73.6%	66.1%	53.6%	43.0%	18-24	
68.7%	70.2%	66.0%	54.9%	75.9%	68.7%	66.5%	59.6%	78.3%	65.4%	64.3%	49.2%	73.1%	66.0%	60.4%	53.5%	25-34	
74.9%	68.4%	64.6%	56.7%	73.8%	71.3%	66.2%	63.5%	76.2%	64.5%	63.7%	56.6%	73.9%	67.3%	61.5%	54.7%	35-44	
76.5%	74.4%	69.9%	63.4%	78.8%	73.5%	70.1%	65.0%	77.5%	73.5%	68.4%	60.4%	78.5%	72.9%	67.8%	60.7%	45-54	
85.1%	83.6%	79.9%	72.2%	83.6%	80.6%	77.6%	76.2%	85.1%	82.0%	78.6%	72.9%	85.1%	81.0%	75.5%	70.2%	55-64	
87.5%	87.1%	86.7%	83.8%	92.1%	90.0%	89.3%	86.8%	92.0%	89.7%	88.6%	85.2%	91.8%	88.9%	86.4%	82.3%	65-74	
77.3%	72.5%	77.2%	65.7%	82.2%	80.8%	80.3%	71.2%	NA*	67.7%	72.2%	60.3%	79.8%	73.2%	64.8%	56.7%	75-84	
80.6%	77.9%	71.7%	68.0%	87.7%	83.2%	80.4%	72.3%	76.9%	77.8%	71.3%	58.8%	82.5%	74.0%	67.1%	62.3%	85+	
75.2%	77.2%	68.7%	66.2%	85.7%	82.0%	79.0%	71.3%	79.3%	74.0%	68.4%	59.4%	81.3%	74.1%	68.1%	62.8%	18-24	
83.1%	82.0%	77.5%	72.0%	84.4%	82.5%	79.0%	76.4%	80.5%	80.4%	76.8%	67.4%	83.3%	79.2%	74.0%	68.7%	25-34	
87.9%	86.5%	83.9%	76.2%	87.7%	86.2%	81.4%	81.4%	88.9%	84.6%	82.7%	77.4%	88.9%	85.1%	80.6%	75.4%	35-44	
87.0%	84.9%	86.0%	83.4%	93.2%	91.8%	90.5%	89.0%	93.0%	89.9%	90.3%	83.8%	92.9%	90.1%	86.9%	83.5%	45-54	

שיעור בעלי תת HbA1c בגובה מ-9% בחולי סוכרת בני 18 ומעלה																← SES	↓ מינ
מאוחדת				מכבי				לאומית				כללית					
8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3		
8.1%	12.0%	16.9%	26.0%	5.5%	10.0%	14.2%	19.2%	NA*	16.7%	17.8%	25.4%	9.6%	14.9%	23.9%	29.8%	18-24	
8.4%	9.8%	10.0%	17.2%	6.3%	10.8%	10.5%	16.9%	3.3%	14.3%	13.5%	19.5%	9.0%	13.6%	17.2%	20.9%	25-34	
8.5%	11.4%	13.3%	16.0%	7.3%	8.3%	11.3%	13.0%	13.1%	13.9%	14.2%	17.2%	7.7%	11.9%	15.5%	19.6%	35-44	
5.7%	8.1%	9.3%	12.3%	5.1%	7.4%	9.3%	11.4%	7.2%	8.6%	10.1%	13.5%	6.6%	8.7%	11.8%	14.9%	45-54	
3.3%	4.1%	5.4%	7.9%	3.5%	4.6%	5.7%	6.3%	3.5%	4.5%	6.1%	9.8%	3.8%	5.3%	7.7%	9.7%	55-64	
2.2%	2.8%	3.9%	5.5%	2.0%	3.1%	4.1%	5.3%	3.7%	4.0%	4.8%	6.4%	2.8%	4.3%	5.6%	7.7%	65-74	
2.2%	3.1%	3.2%	4.9%	3.1%	3.7%	4.1%	6.7%	6.0%	5.0%	4.3%	5.0%	3.7%	5.0%	5.2%	6.1%	75-84	
6.1%	9.4%	12.6%	16.8%	4.0%	7.7%	7.0%	10.7%	NA*	15.4%	19.1%	18.3%	6.5%	10.5%	17.1%	21.0%	85+	
6.0%	4.9%	10.7%	11.8%	4.1%	5.5%	5.6%	10.4%	7.7%	9.4%	11.2%	23.2%	6.5%	8.3%	13.3%	16.1%	18-24	
8.9%	8.4%	10.3%	10.2%	3.3%	5.6%	8.3%	8.9%	8.1%	11.0%	12.4%	16.9%	6.5%	9.9%	13.5%	15.3%	25-34	
5.2%	4.7%	7.0%	8.7%	3.6%	4.8%	6.1%	8.1%	5.8%	6.8%	8.1%	12.4%	5.2%	6.9%	9.7%	12.3%	35-44	
2.8%	3.1%	4.0%	5.7%	2.4%	3.1%	4.4%	5.4%	3.3%	4.5%	5.5%	8.2%	2.8%	4.2%	6.2%	8.5%	45-54	
2.2%	3.3%	3.4%	4.9%	2.2%	2.9%	3.7%	4.4%	2.4%	3.6%	4.0%	7.2%	2.4%	3.6%	5.3%	6.9%	55-64	
1.4%	3.5%	3.2%	1.9%	3.9%	3.6%	4.3%	4.7%	3.0%	3.1%	3.3%	5.6%	3.4%	4.0%	5.4%	6.4%	65-74	

שיעור הביצוע של בדיקות עיניים בחולי סוכרת בני 18-84																← SES	↓ מינ
מאוחדת				מכבי				לאומית				כללית					
8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3		
46.9%	47.0%	62.8%	59.9%	60.5%	57.9%	58.5%	56.6%	NA*	36.0%	48.6%	42.2%	65.2%	64.9%	64.8%	67.5%	18-24	
52.3%	57.6%	57.0%	61.9%	60.1%	60.5%	64.8%	68.0%	34.7%	45.8%	37.9%	41.5%	62.5%	65.3%	69.9%	69.7%	25-34	
63.9%	58.5%	59.6%	65.5%	68.9%	70.4%	71.6%	74.7%	38.0%	46.2%	47.7%	53.8%	68.3%	69.0%	72.2%	73.0%	35-44	
64.0%	62.9%	65.0%	67.2%	76.1%	77.4%	77.9%	77.0%	55.4%	50.4%	51.7%	54.5%	71.3%	71.4%	75.7%	75.7%	45-54	
71.0%	69.0%	68.7%	71.9%	83.1%	83.7%	81.8%	81.5%	61.3%	60.9%	58.5%	60.8%	78.1%	78.0%	78.7%	79.0%	55-64	
72.5%	71.2%	72.1%	65.9%	84.2%	83.3%	80.0%	79.4%	65.7%	63.0%	60.0%	64.1%	78.8%	78.2%	76.3%	75.3%	65-74	
57.3%	54.1%	57.5%	64.5%	49.7%	43.7%	43.3%	53.5%	47.5%	50.0%	45.3%	52.5%	66.7%	66.9%	68.7%	71.8%	75-84	
58.5%	60.6%	60.0%	63.1%	48.2%	50.5%	52.1%	58.9%	47.1%	47.6%	44.8%	47.8%	62.6%	67.7%	70.2%	75.3%	85+	
64.4%	65.8%	68.0%	71.5%	61.7%	62.7%	66.2%	71.1%	50.0%	52.2%	57.3%	61.8%	70.0%	73.9%	76.9%	79.0%	18-24	
70.8%	65.6%	69.9%	74.3%	75.2%	75.9%	75.5%	77.4%	58.6%	57.1%	60.8%	64						

סוכרת																קבוצת גיל ↓	מין ↓	
מאוחדת				מבוי				לאומית				כללית						
8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3			
שיעור התישוד של GFR בחולי סוכרת בני 84-18																גברים	81.6%	18-34
82.7%	79.8%	81.6%	79.6%	78.3%	74.7%	76.0%	80.0%	NA*	67.4%	73.7%	69.0%	78.7%	78.2%	80.4%	81.6%		18-34	
74.3%	79.9%	81.7%	84.3%	78.7%	77.4%	80.3%	84.0%	80.6%	71.1%	73.4%	70.4%	79.6%	79.3%	80.9%	82.2%		35-44	
82.5%	80.2%	83.3%	85.4%	83.4%	85.3%	87.1%	88.1%	76.7%	76.5%	80.2%	81.3%	83.7%	84.5%	84.4%	87.0%		45-54	
86.2%	86.9%	86.7%	88.7%	89.4%	90.4%	90.7%	91.6%	82.7%	83.2%	84.1%	84.9%	88.9%	88.7%	88.8%	89.7%		55-64	
90.9%	91.5%	91.8%	93.1%	94.6%	95.3%	94.8%	95.1%	90.5%	88.8%	89.9%	88.4%	93.4%	93.9%	93.3%	93.3%		65-74	
94.6%	95.5%	94.7%	94.7%	96.7%	97.0%	96.5%	96.8%	92.3%	92.8%	91.3%	91.6%	96.0%	95.7%	95.2%	94.3%		75-84	
80.5%	83.7%	86.1%	89.4%	78.3%	75.9%	77.0%	82.2%	75.0%	74.5%	78.7%	87.1%	87.5%	84.3%	86.8%	88.4%		18-34	
82.9%	83.7%	83.3%	88.5%	74.8%	77.8%	77.6%	82.9%	70.6%	81.4%	76.3%	78.8%	83.2%	84.2%	87.1%	88.8%		35-44	
87.3%	86.9%	88.8%	92.8%	79.4%	82.1%	86.6%	88.8%	91.3%	83.4%	87.5%	86.7%	87.1%	88.3%	89.4%	91.0%		45-54	
87.1%	87.2%	89.9%	93.5%	87.7%	89.2%	90.8%	93.5%	84.5%	85.4%	88.2%	90.4%	90.2%	91.3%	91.5%	93.5%	55-64		
93.8%	93.3%	93.3%	94.8%	93.8%	94.8%	95.7%	96.4%	90.9%	91.5%	90.9%	89.6%	94.1%	94.0%	94.2%	94.9%	65-74		
94.4%	93.7%	94.1%	94.3%	96.1%	96.3%	96.5%	96.1%	90.5%	91.5%	90.7%	90.7%	95.7%	95.3%	95.2%	95.3%	75-84		
שיעור חולי סוכרת עם פגיעה בליתיות בני 84-18																גברים	20.3%	18-34
6.1%	9.9%	14.0%	16.1%	5.1%	7.3%	7.0%	14.1%	NA*	9.9%	16.3%	19.6%	7.7%	12.2%	15.9%	20.3%		18-34	
18.9%	16.5%	17.6%	25.8%	10.6%	13.5%	13.0%	19.5%	18.6%	18.5%	22.3%	23.1%	12.9%	18.6%	23.0%	24.6%		35-44	
17.8%	23.8%	26.3%	25.1%	13.7%	16.0%	18.8%	22.4%	26.8%	21.4%	25.1%	28.2%	16.6%	22.4%	25.6%	27.3%		45-54	
24.7%	27.7%	30.0%	32.1%	20.2%	22.6%	26.4%	28.8%	28.6%	26.9%	27.4%	32.6%	23.3%	28.4%	31.0%	33.0%		55-64	
37.7%	41.0%	39.5%	44.3%	32.9%	34.9%	35.2%	36.3%	34.3%	31.7%	33.4%	38.0%	34.7%	38.2%	41.0%	43.3%		65-74	
53.5%	54.5%	56.8%	57.3%	51.8%	53.9%	53.9%	50.2%	43.1%	41.3%	39.3%	43.7%	52.1%	52.4%	53.9%	55.8%		75-84	
9.1%	12.9%	17.2%	19.3%	4.3%	3.2%	5.3%	8.7%	24.2%	7.3%	13.0%	22.5%	8.1%	13.7%	14.5%	20.8%		18-34	
18.7%	13.8%	15.7%	18.4%	5.4%	7.4%	8.7%	10.3%	19.0%	12.4%	20.8%	23.7%	13.8%	14.7%	18.2%	20.1%		35-44	
16.0%	18.7%	21.4%	21.3%	7.5%	9.5%	12.1%	14.2%	15.6%	12.6%	17.7%	21.8%	14.3%	15.8%	19.0%	23.1%		45-54	
15.5%	20.3%	21.5%	24.8%	11.1%	13.7%	16.4%	19.8%	20.3%	18.8%	19.6%	23.6%	16.2%	20.1%	23.7%	25.2%	55-64		
27.7%	28.4%	33.8%	34.8%	22.4%	25.8%	26.9%	27.7%	19.3%	23.0%	23.0%	25.3%	25.4%	28.7%	32.6%	35.4%	65-74		
44.7%	49.1%	52.0%	53.7%	40.7%	45.9%	46.5%	47.7%	25.8%	27.6%	26.3%	31.7%	44.8%	47.0%	50.8%	54.1%	75-84		
שיעור התישוד של מדידות לוחץ דם בחולי סוכרת בני 18 ומעלה																גברים	78.0%	18-34
51.9%	57.4%	69.2%	74.9%	63.8%	62.1%	61.1%	66.2%	NA*	43.0%	51.4%	62.6%	59.5%	69.1%	69.7%	78.0%		18-34	
71.6%	68.7%	71.7%	79.6%	65.7%	66.1%	68.6%	75.2%	54.2%	59.5%	60.4%	65.7%	69.0%	71.9%	76.2%	81.8%		35-44	
71.8%	71.6%	77.5%	82.0%	71.5%	74.7%	78.3%	81.2%	60.2%	64.7%	68.1%	74.1%	75.3%	79.0%	82.4%	86.2%		45-54	
77.9%	78.8%	83.8%	86.3%	79.7%	82.7%	82.9%	86.2%	71.8%	76.0%	76.4%	79.6%	83.2%	85.7%	87.9%	89.8%		55-64	
86.2%	87.1%	86.4%	90.3%	87.4%	89.0%	88.3%	89.9%	81.3%	82.5%	83.2%	88.9%	88.9%	90.8%	91.6%	93.0%		65-74	
91.1%	89.7%	89.6%	90.0%	89.8%	91.7%	91.9%	90.7%	85.1%	83.4%	81.1%	86.5%	92.1%	93.4%	93.6%	93.3%		75-84	
83.9%	89.9%	86.6%	87.2%	87.5%	89.4%	88.1%	86.5%	86.9%	79.8%	76.9%	74.1%	92.0%	93.1%	92.9%	92.0%		85+	
73.2%	71.5%	78.6%	80.4%	59.7%	60.3%	60.4%	65.4%	52.5%	56.4%	67.3%	71.2%	66.3%	68.7%	77.1%	81.2%		18-34	
81.7%	70.0%	73.3%	83.7%	60.6%	62.1%	64.8%	71.0%	54.9%	70.3%	69.4%	68.1%	68.1%	70.6%	76.8%	85.0%		35-44	
73.9%	76.4%	80.7%	86.8%	65.0%	69.0%	74.7%	79.7%	68.7%	69.9%	79.2%	78.7%	75.5%	80.9%	84.1%	87.9%	45-54		
78.7%	79.2%	82.6%	88.7%	75.0%	79.9%	82.4%	85.5%	75.4%	78.1%	80.8%	83.8%	84.5%	86.6%	89.3%	92.4%	55-64		
86.5%	86.4%	87.6%	91.4%	86.8%	88.5%	88.9%	90.2%	83.2%	82.8%	83.6%	84.0%	90.2%	91.2%	91.7%	93.3%	65-74		
88.0%	87.8%	87.4%	88.9%	88.9%	89.9%	88.9%	89.1%	82.4%	82.7%	80.8%	78.9%	91.3%	92.3%	92.4%	93.2%	75-84		
81.0%	81.3%	80.3%	79.9%	79.4%	82.0%	83.2%	82.0%	64.0%	73.2%	69.9%	71.1%	85.6%	89.5%	88.7%	91.0%	85+		
שיעור בעלי לוחץ דם נמוך או שווה ל-140/90 מ"מ בחולי סוכרת בני 18 ומעלה																גברים	92.8%	18-34
97.6%	91.4%	95.7%	90.8%	92.4%	93.4%	91.7%	91.9%	NA*	89.2%	88.9%	92.3%	93.1%	94.9%	93.6%	92.8%		18-34	
83.3%	83.2%	84.3%	87.6%	83.8%	83.3%	82.3%	84.5%	84.6%	87.6%	86.8%	88.2%	87.1%	85.3%	84.1%	87.7%		35-44	
81.7%	79.0%	78.5%	81.4%	82.3%	81.0%	80.3%	82.0%	78.8%	76.7%	82.3%	85.7%	82.9%	81.2%	82.3%	84.0%		45-54	
78.6%	76.1%	76.5%	78.4%	80.5%	78.8%	79.2%	80.9%	78.2%	78.0%	80.3%	82.8%	78.8%	78.5%	77.8%	81.2%		55-64	
76.0%	74.6%	74.4%	76.8%	79.9%	78.7%	78.6%	79.4%	80.2%	80.1%	80.4%	81.0%	80.0%	77.4%	76.2%	79.6%		65-74	
75.2%	73.0%	72.5%	74.1%	77.6%	77.0%	78.4%	77.7%	82.2%	79.6%	80.0%	83.2%	79.6%	76.1%	75.9%	79.3%		75-84	
72.9%	78.5%	79.1%	75.5%	75.3%	76.9%	78.7%	78.4%	78.8%	83.0%	84.0%	77.7%	77.5%	75.6%	75.3%	77.0%		85+	
98.3%	96.7%	95.5%	96.9%	95.5%	93.1%	94.2%	90.7%	NA*	94.3%	93.1%	94.9%	97.4%	95.5%	95.8%	96.4%		18-34	
91.0%	89.4%	89.5%	94.9%	89.6%	90.7%	90.2%	91.4%	NA*	93.1%	93.2%	93.5%	92.2%	90.4%	90.8%	90.7%		35-44	
86.7%	82.8%	81.1%	83.1%	87.3%	85.1%	85.9%	85.5%	87.4%	86.0%	86.7%	87.1%	87.4%	85.7%	84.6%	86.3%	45-54		
81.5%	79.8%	80.8%	81.7%	84.3%	82.8%	83.4%	83.9%	82.8%	80.6%	81.8%	82.6%	84.8%	82.3%	81.4%	83.0%	55-64		
76.2%	76.0%	75.1%	77.2%	79.8%	79.4%	81.3%	79.4%	80.3%	79.8%	78.8%	82.3%	80.5%	78.7%	78.2%	78.4%	65-74		
69.0%	69.7%	71.3%	71.5%	75.0%	74.1%	75.8%	72.8%	78.9%	79.4%	76.7%	80.3%	74.6%	73.5%	73.5%	76.1%	75-84		
73.5%	63.0%	66.3%	68.1%	68.4%	68.5%	73.7%	72.0%	75.0%	76.9%	77.4%	76.7%	71.4%	70.1%	72.6%	77.1%	85+		
שיעור הבעיות של בדיקת כולסטרול בחולי סוכרת בני 18 ומעלה																גברים	78.1%	18-34
82.7%	75.4%	82.3%	78.5%	72.1%	68.3%	70.0%	75.2%	NA*	67.4%	72.6%	66.3%	76.9%	75.8%	78.2%	78.1%		18-34	
76.1%	79.5%	81.5%	83.9%	73.9%	74.0%	76.3%	80.4%	79.2%	68.9%	70.4%	70.1%	77.9%	77.3%	79.2%	80.3%		35-44	
82.7%	79.4%	82.5%	85.1%	80.0%	82.2%	83.7%	85.6%	75.9%	75.2%	77.9%	81.2%	82.5%	83.2%	83.2%	85.6%		45-54	
85.6%	86.1%	86.4%	88.2%	87.0%	88.2%	88.3%	89.5%	81.3%	83.0%	83.0%	84.0%	87.5%	87.2%	87.4%	88.5%		55-64	
90.0%	90.8%	90.9%	91.9%	92.7%	93.1%	92.8%	92.9%	92.7%	88.0%	88.9%	88.1%	92.1%	92.6%	92.2%	92.1%		65-74	
93.4%	94.5%	93.7%	93.2%	94.6%	95.1%	94.0%	93.9%	90.8%	90.9%	89.0%	89.1%	94.3%	94.2%	93.5%	92.9%		75-84	
92.2%	91.3%	90.7%	92.1%	91.8%	90.9%	92.2%	92.0%	82.3%	84.4%	86.0%	87.1%	92.4%	92.3%	92.6%	91.9%		85+	
79.3%	80.8%	81.7%	86.9%	64.2%	58.1%	57.6%	67.1%	72.5%	69.1%	75.3%	84.9%	83.2%	78.7%	82.8%	84.2%		18-34	
78.0%	82.3%	80.7%	87.3%	58.9%	63.1%	63.8%	71.7%	70.6%	80.0%	72.0%	74.8%	78.2%	80.9%	85.1%	86.3%		35-44	
87.3%	85.7%	88.5%	92.5%	70.1%	71.9%	78.6%	82.9%	88.7%	83.7%	87.4%	87.0%	85.7%	87.1%	88.1%	90.0%	45-54		
87.5%	86.4%	89.2%	92.9%	82.1%	84.6%	86.8%	91.1%	85.6%	84.7%	87.6%	89.8%	89.0%	90.0%	90.4%	92.5%	55-64		
93.4%	92.3%	92.6%	94.2%	91.5%	92.3%	93.2%	93.5%	91.0%	90.8%	90.7%	88.8%	93.1%	93.2%	93.0%	93.8%	65-74		
92.8%	91.9%	92.7%	93.2%	93.2%	93.9%	93.6%	93.3%	88.5%	89.3%	88.6%	88.2%	93.7%	93.6%	93.4%	93.5%	75-84		
88.8%	88.4%	90.2%	88.3%	87.2%	89.6%	89.4%	91.0%	82.0%	85.1%	79.4%	81.5%	90.5%	90.5%	90.8%	91.			

סוכרת																
מאוחדת				מכבי				לאומית				כללית				← SES קבוצת גיל ↓ מין
8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	
שיעור היתעד של GFR בחולי סוכרת בני 18-84																
שיעור ההמצאות של עודף משקל בקרב חולי סוכרת בני 64-20																
35.6%	27.4%	27.7%	24.4%	30.7%	26.4%	29.6%	24.0%	NA*	28.2%	32.6%	23.1%	34.9%	32.0%	28.5%	28.2%	20-34
31.7%	32.2%	32.2%	30.6%	33.2%	29.7%	27.3%	27.2%	39.1%	31.7%	38.9%	29.5%	39.4%	33.5%	33.2%	33.6%	35-44
39.8%	35.2%	36.6%	35.5%	39.4%	36.2%	36.5%	33.3%	39.9%	36.2%	36.0%	39.6%	40.6%	38.2%	36.4%	38.5%	45-54
47.7%	41.2%	40.1%	41.4%	45.0%	41.8%	39.7%	38.0%	46.3%	42.8%	41.5%	41.8%	45.4%	44.3%	42.3%	42.9%	55-64
23.7%	26.2%	28.5%	26.0%	31.4%	23.8%	21.3%	23.6%	NA*	30.4%	23.1%	26.0%	24.4%	28.4%	24.2%	21.8%	20-34
32.3%	26.9%	25.1%	26.8%	28.9%	26.2%	25.8%	23.7%	18.2%	21.4%	23.5%	19.3%	26.7%	26.4%	27.3%	25.9%	35-44
28.3%	31.8%	27.1%	24.7%	32.7%	31.4%	28.6%	24.7%	32.7%	25.2%	24.2%	26.8%	29.3%	31.2%	29.8%	25.0%	45-54
39.4%	35.8%	32.5%	26.5%	37.7%	32.4%	30.9%	27.9%	32.5%	37.5%	33.5%	28.6%	37.1%	33.9%	31.4%	27.5%	55-64
שיעור ההמצאות של השמנת יתר בקרב חולי סוכרת בני 64-20																
11.1%	17.0%	17.0%	22.5%	15.5%	20.2%	21.5%	24.0%	NA*	33.3%	22.8%	27.8%	14.1%	15.6%	18.8%	22.1%	20-34
28.0%	27.5%	26.1%	33.6%	28.6%	32.0%	31.1%	33.1%	26.1%	31.7%	26.0%	32.6%	24.9%	29.5%	29.4%	31.8%	35-44
29.5%	35.1%	30.4%	33.7%	30.8%	32.0%	32.2%	33.9%	32.9%	34.8%	31.0%	31.5%	27.9%	30.1%	31.1%	31.7%	45-54
23.7%	31.0%	31.0%	31.4%	29.3%	30.6%	31.6%	31.7%	22.1%	27.0%	29.0%	29.4%	27.2%	27.2%	27.3%	27.8%	55-64
13.6%	25.4%	19.4%	21.1%	20.8%	26.8%	28.3%	24.0%	NA*	16.1%	29.7%	22.1%	17.5%	17.5%	22.4%	23.6%	20-34
14.5%	28.7%	29.2%	30.4%	30.1%	29.7%	30.3%	30.1%	30.3%	26.2%	24.7%	28.6%	22.9%	28.5%	28.1%	31.8%	35-44
27.0%	27.7%	31.0%	34.5%	32.0%	32.2%	32.5%	32.9%	28.8%	35.0%	34.2%	33.7%	25.1%	28.0%	30.8%	33.3%	45-54
24.5%	28.4%	30.7%	35.3%	28.4%	31.9%	31.8%	33.4%	32.1%	26.0%	31.0%	32.3%	26.4%	29.0%	30.5%	32.6%	55-64
שיעור ההמצאות של השמנת יתר חלנית בקרב חולי סוכרת בני 64-20																
8.9%	12.3%	11.2%	24.4%	20.1%	22.8%	22.1%	23.5%	NA*	10.3%	21.7%	21.3%	9.2%	12.9%	16.9%	18.9%	20-34
19.5%	25.6%	24.7%	25.0%	23.3%	24.1%	30.3%	29.0%	13.0%	21.4%	21.2%	26.8%	17.3%	20.0%	20.7%	20.2%	35-44
15.9%	17.3%	21.1%	19.8%	16.7%	21.1%	21.0%	24.1%	16.2%	14.4%	20.7%	18.9%	15.8%	17.3%	17.6%	16.5%	45-54
10.8%	13.2%	16.1%	14.1%	10.4%	14.3%	15.7%	16.6%	10.2%	13.1%	12.7%	14.8%	9.8%	11.0%	11.7%	12.5%	55-64
16.9%	20.5%	24.7%	24.4%	21.8%	29.7%	32.6%	33.2%	NA*	26.8%	24.2%	26.9%	17.2%	22.3%	25.8%	26.2%	20-34
35.5%	26.3%	33.7%	32.5%	24.7%	30.0%	31.7%	34.9%	36.4%	36.9%	39.2%	46.6%	21.8%	27.2%	30.3%	33.2%	35-44
27.0%	25.5%	31.0%	32.5%	20.8%	24.9%	29.3%	34.1%	24.0%	27.7%	29.9%	33.7%	25.8%	25.1%	27.4%	33.9%	45-54
16.5%	21.2%	26.7%	30.1%	17.9%	21.8%	26.7%	30.3%	17.0%	21.9%	25.2%	30.1%	16.8%	20.4%	24.8%	30.8%	55-64
שיעור היתעד של מרכיבי BMI בחולי סוכרת בני 64-20																
81.8%	84.2%	82.0%	84.6%	72.6%	73.9%	73.6%	79.8%	77.4%	76.4%	79.6%	76.6%	83.5%	85.1%	86.3%	87.7%	65-69
83.9%	84.6%	83.7%	87.3%	74.6%	78.8%	77.5%	80.2%	84.1%	83.9%	81.2%	84.1%	89.2%	89.4%	89.4%	91.2%	70-74
86.4%	86.0%	84.8%	86.1%	76.3%	80.4%	79.5%	81.0%	80.0%	80.0%	80.0%	82.8%	89.5%	90.6%	89.5%	90.1%	75-79
83.2%	84.3%	81.3%	80.2%	74.0%	78.2%	76.6%	78.9%	79.9%	73.0%	69.8%	76.4%	89.7%	88.6%	87.5%	88.3%	80-84
82.2%	82.7%	84.1%	88.6%	71.9%	76.5%	76.8%	78.0%	78.7%	78.1%	81.9%	82.0%	85.9%	86.1%	87.5%	88.8%	65-69
87.6%	84.3%	83.1%	87.3%	74.1%	77.4%	77.4%	79.1%	83.4%	83.1%	82.8%	81.6%	89.6%	89.8%	88.8%	90.2%	70-74
83.7%	83.0%	80.5%	81.7%	74.4%	76.2%	74.9%	75.8%	79.3%	77.7%	75.5%	75.0%	89.7%	89.2%	88.5%	87.8%	75-79
77.5%	76.6%	76.2%	73.3%	73.1%	70.4%	73.1%	75.3%	71.1%	69.0%	67.9%	65.5%	86.0%	86.0%	85.1%	84.1%	80-84
שיעור ההמצאות של השמנת יתר והשמנת יתר חלנית בקרב חולי סוכרת בני 64-20																
34.7%	38.6%	39.1%	39.7%	32.4%	38.4%	41.8%	42.0%	35.0%	33.3%	36.7%	38.6%	32.3%	33.2%	34.7%	36.9%	65-69
29.7%	37.0%	40.8%	42.7%	29.0%	34.6%	39.8%	42.1%	26.9%	33.9%	33.7%	38.0%	28.7%	30.8%	31.9%	33.9%	70-74
28.8%	29.7%	36.8%	36.4%	26.3%	31.0%	36.0%	35.1%	25.5%	28.4%	32.6%	31.2%	24.9%	27.9%	30.2%	31.9%	75-79
21.8%	27.3%	32.0%	37.1%	20.9%	26.5%	27.9%	28.1%	18.7%	21.5%	28.5%	33.8%	20.9%	22.6%	26.2%	29.9%	80-84
42.2%	47.3%	56.1%	62.7%	39.1%	48.3%	55.4%	63.4%	33.3%	41.5%	55.0%	57.2%	38.3%	44.0%	51.5%	61.0%	65-69
35.3%	44.9%	51.5%	60.1%	37.8%	46.8%	53.9%	57.8%	37.5%	43.2%	51.8%	58.4%	36.2%	41.4%	49.0%	56.8%	70-74
31.4%	44.0%	50.2%	56.5%	35.2%	44.7%	49.9%	55.7%	32.2%	40.2%	49.4%	52.1%	33.1%	39.5%	45.8%	52.8%	75-79
29.0%	39.8%	44.4%	47.9%	32.1%	40.9%	46.3%	51.2%	27.4%	38.9%	43.0%	48.4%	29.7%	36.7%	41.3%	48.6%	80-84
שיעור קבלת חיסון נגד שפעת בחולי סוכרת בני 18 ומעלה																
30.9%	19.7%	21.8%	23.5%	24.4%	14.5%	13.4%	14.1%	NA*	40.7%	37.7%	45.5%	22.7%	22.0%	21.8%	35.4%	18-34
33.0%	25.9%	24.9%	25.9%	22.1%	16.2%	16.4%	15.8%	43.1%	48.9%	46.2%	49.0%	27.0%	22.2%	27.4%	38.5%	35-44
37.0%	30.6%	29.8%	32.9%	25.2%	21.6%	20.6%	25.0%	45.5%	49.5%	54.4%	61.7%	33.0%	31.2%	36.2%	44.8%	45-54
47.3%	42.5%	39.3%	41.8%	41.0%	31.8%	29.6%	30.3%	61.0%	61.2%	62.6%	65.7%	47.6%	45.6%	46.6%	53.2%	55-64
72.7%	63.4%	56.4%	55.3%	59.5%	49.7%	43.5%	41.2%	77.1%	75.3%	74.7%	74.8%	69.5%	63.4%	61.1%	64.0%	65-74
81.5%	73.5%	65.3%	58.6%	68.6%	58.4%	49.1%	48.4%	84.8%	79.9%	78.4%	79.1%	79.6%	74.2%	70.0%	69.7%	75-84
78.8%	74.9%	68.5%	64.6%	67.5%	57.5%	49.7%	44.3%	83.8%	78.8%	76.0%	79.1%	82.8%	75.2%	72.4%	72.1%	85+
35.4%	25.6%	26.6%	32.7%	23.0%	16.7%	14.5%	17.0%	42.5%	46.8%	54.0%	59.7%	29.9%	22.9%	24.3%	35.9%	18-34
26.8%	28.1%	27.3%	26.2%	23.7%	16.6%	11.4%	11.3%	45.1%	52.4%	55.2%	58.0%	32.0%	23.9%	28.0%	39.4%	35-44
36.6%	27.9%	30.4%	34.6%	23.4%	18.9%	18.1%	17.9%	60.7%	52.7%	67.9%	64.0%	32.8%	31.8%	35.7%	45.5%	45-54
49.2%	39.1%	36.5%	40.9%	37.8%	30.3%	26.2%	25.5%	64.4%	63.7%	71.4%	67.0%	46.7%	41.9%	41.8%	51.7%	55-64
68.4%	58.4%	50.7%	48.5%	58.3%	45.4%	37.8%	37.5%	79.6%	72.8%	74.7%	74.8%	66.0%	57.8%	53.1%	57.4%	65-74
76.7%	65.8%	58.2%	52.3%	61.1%	50.0%	43.3%	38.2%	82.6%	74.8%	73.3%	74.0%	73.6%	65.3%	61.3%	62.4%	75-84
71.1%	64.7%	56.8%	56.1%	52.7%	41.5%	40.8%	40.6%	72.0%	71.9%	70.4%	74.4%	75.5%	68.0%	65.1%	65.0%	85+
שיעור קבלת חיסון נגד פנואמוניק בחולי סוכרת בני 74-65																
74.0%	70.2%	69.2%	76.6%	74.4%	71.9%	71.0%	73.9%	92.0%	91.0%	90.1%	89.2%	73.8%	72.9%	74.5%	78.9%	65-69
89.9%	89.6%	87.7%	89.1%	92.1%	91.2%	90.6%	89.8%	98.5%	98.4%	96.7%	97.6%	90.1%	89.6%	88.5%	91.9%	70-74
73.1%	65.9%	65.7%	73.2%	74.4%	71.0%	67.4%	69.9%	95.0%	92.2%	89.9%	87.8%	74.5%	70.3%	70.3%	76.0%	65-69
87.0%	83.9%	84.1%	85.4%	89.9%	87.4%	87.4%	87.6%	98.9%	98.9%	98.2%	97.1%	87.7%	85.6%	85.2%	89.3%	70-74
שיעור ההמצאות של סוכרת בקרב בני 17-2																
0.0%	0.1%	0.0%	0.0%	0.0%	0.1%	0.0%	0.1%	0.2%	0.1%	0.1%	0.0%	0.0%	0.1%	0.0%	0.1%	2-9
0.2%	0.2%	0.3%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	10-17
0.0%	0.0%	0.0%	0.1%	0.0%	0.0%	0.0%	0.1%	0.0%	0.1%	0.0%	0.1%	0.1%	0.1%	0.0%	0.1%	2-9
0.3%	0.2%	0.2%	0.3%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	0.2%	0.3%	10-17
שיעור הנבדקים במרפאת סוכרת ילדים בקרב חולי סוכרת בני 17-2																
NA*	NA*	NA*	NA*	NA*	90.9%	NA*	NA*	NA*	NA*	NA*	NA*	NA*	87.3%	90.7%	83.9%	2-9
NA*	73.3%	54.3%	52.1%	91.6%	90.2%	94.7%	83.7%	NA*	NA*	76.2%	79.5%	92.8%	89.6%	84.9%	81.1%	10-17
NA*	NA*	NA*	NA*	NA*	NA*	NA*	NA*	NA*	NA*	NA*	NA*	93.8%	93.5%	79.1%	86.2%	2-9
NA*	65.7%	52.1%	47.8%	94.4%	89.3%	96.5%	78.0%	NA*	NA*	75.8%	73.7%	89.9%	91.4%	83.7%	80.0%	10-17

חא מסומן ב-* יעוד עור מכנה נמוך מ 30 מבוטחים

מחלות זיהומיות															
מאוחדת				מכבי				לאומית				כללית			
8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3
שיעור השימוש בתרופות אנטיביוטיות מסוג צפולספורינים וקווינלונים, מתוך סך התרופות האנטיביוטיות															
<div style="display: flex; justify-content: space-between;"> ← SES ↓ מין </div> <div style="display: flex; justify-content: space-between;"> קבוצת גיל ↓ </div>															
6.7%	6.8%	7.4%	6.5%	12.9%	11.7%	11.3%	10.3%	8.0%	7.0%	7.6%	6.6%	11.0%	9.4%	8.0%	5.8%
11.3%	11.6%	13.4%	14.3%	13.0%	12.8%	13.1%	14.6%	7.4%	10.6%	11.8%	12.8%	12.7%	12.3%	12.9%	14.3%
21.3%	19.7%	19.7%	19.7%	18.4%	15.7%	14.9%	16.3%	14.7%	14.7%	15.0%	15.9%	19.5%	18.1%	16.8%	18.5%
20.1%	22.1%	21.2%	22.0%	16.8%	16.7%	17.0%	18.2%	17.7%	17.0%	15.1%	19.3%	18.1%	18.1%	17.6%	18.7%
24.8%	26.1%	26.1%	28.5%	21.0%	19.5%	20.7%	20.4%	21.2%	19.4%	20.2%	23.3%	20.2%	21.3%	20.6%	21.6%
30.1%	31.3%	33.8%	34.1%	24.1%	24.0%	24.1%	24.4%	23.0%	26.8%	23.9%	27.4%	24.5%	25.1%	25.0%	26.5%
35.5%	36.9%	36.6%	36.2%	29.6%	28.8%	30.2%	30.4%	29.7%	29.6%	30.7%	28.2%	29.0%	29.8%	30.4%	31.0%
37.0%	41.3%	41.4%	44.9%	33.6%	34.4%	35.9%	33.5%	34.6%	31.5%	34.2%	35.6%	33.1%	35.5%	36.4%	36.4%
40.8%	40.4%	46.7%	45.7%	40.0%	39.0%	39.4%	42.0%	39.0%	37.7%	38.3%	37.9%	38.0%	39.4%	42.2%	41.6%
43.6%	46.8%	43.0%	45.9%	42.7%	44.8%	44.0%	42.8%	42.5%	51.1%	42.5%	45.5%	43.4%	46.6%	46.3%	46.6%
9.1%	8.7%	9.3%	9.1%	15.7%	15.0%	15.0%	12.7%	10.4%	9.0%	8.7%	8.9%	13.7%	12.0%	10.7%	9.0%
15.8%	13.6%	13.4%	13.0%	16.4%	16.2%	14.9%	13.5%	13.3%	12.0%	12.0%	12.0%	16.8%	16.2%	15.0%	16.0%
29.0%	26.7%	27.8%	27.4%	20.9%	20.2%	20.7%	21.4%	24.8%	21.6%	22.1%	22.9%	24.9%	24.9%	24.6%	26.6%
27.0%	29.7%	29.0%	29.7%	21.3%	20.6%	20.6%	22.2%	25.7%	22.2%	23.4%	23.6%	24.8%	25.5%	25.5%	26.1%
30.5%	32.0%	31.6%	31.1%	23.0%	21.9%	21.3%	22.9%	26.6%	24.7%	25.6%	25.9%	27.1%	27.4%	26.6%	27.0%
34.7%	34.2%	33.0%	33.6%	25.0%	24.2%	24.6%	25.2%	31.8%	28.4%	26.9%	27.6%	29.4%	28.6%	28.1%	28.8%
34.8%	34.9%	35.8%	36.2%	27.1%	27.3%	26.9%	27.1%	31.3%	30.0%	31.9%	32.6%	30.1%	30.8%	31.4%	32.0%
31.7%	34.8%	36.3%	39.4%	29.9%	31.5%	32.5%	31.8%	37.8%	34.9%	38.1%	38.4%	31.1%	33.2%	35.9%	37.1%
32.9%	39.2%	36.1%	45.8%	33.2%	36.3%	38.5%	37.4%	46.3%	42.1%	42.2%	38.3%	33.5%	35.4%	40.1%	40.6%
34.8%	37.1%	39.3%	43.2%	40.4%	41.8%	41.6%	38.7%	49.7%	51.8%	46.0%	49.0%	34.6%	39.3%	42.4%	45.1%
היקף השימוש בתרופות אנטיביוטיות סיסטמיות, ורישוח ל-1000 איש לשנה															
913.5	972.3	898.2	829.9	874.9	874.6	821.3	846.9	1070.9	1085.8	958.8	869.1	820.5	896.1	875.0	879.7
416.9	464.1	463.4	417.5	383.5	378.2	366.8	413.9	426.4	470.2	457.9	441.8	354.7	370.4	370.4	376.3
376.3	427.5	463.8	446.6	382.5	408.4	414.4	449.7	359.0	410.9	421.2	425.0	343.7	395.0	410.7	408.3
419.5	487.5	535.6	561.6	459.0	485.7	483.7	533.9	454.4	461.8	473.7	536.5	416.1	452.6	494.7	550.2
436.6	486.8	571.5	654.9	460.4	476.6	496.1	598.3	448.7	509.6	564.3	657.2	427.9	484.0	547.5	670.7
484.9	590.1	711.6	865.9	521.9	608.3	718.8		530.9	608.0	735.2	856.3	509.1	596.7	723.6	862.9
736.0	790.2	944.3	1183.4	742.4	742.6	753.3	950.5	749.5	846.9	939.7	1079.6	772.5	813.6	899.6	1074.4
950.7	946.8	1130.6	1499.4	924.2	1095.0	881.5	1041.9	998.8	1012.9	1053.9	1337.9	961.5	970.9	1032.7	1373.1
1039.2	1145.7	1323.7	1650.2	1040.0	1040.9	952.0	1073.3	1179.7	1136.3	1250.0	1252.1	1108.0	1121.6	1180.1	1478.4
1229.2	1178.0	1416.0	1701.5	1111.5	1018.6	992.1	1126.6	1193.3	1186.0	1254.7	1461.9	1208.5	1172.8	1210.0	1354.2
937.9	1014.3	908.2	834.4	933.2	905.6	829.1	822.4	1077.1	1110.4	962.4	874.1	845.4	918.8	883.0	831.4
499.9	530.9	524.3	462.2	460.4	469.3	450.3	462.2	534.6	532.5	523.6	501.6	426.2	462.2	434.4	408.7
741.0	820.5	827.5	739.3	743.2	789.8	799.8	770.0	756.7	812.1	784.5	777.8	692.4	774.6	744.9	685.5
815.9	895.9	940.0	859.7	822.5	850.5	840.8	830.6	822.6	922.9	884.5	918.9	763.7	840.2	876.2	879.4
752.3	870.8	920.7	912.3	801.7	837.8	845.6	875.6	778.8	898.6	942.2	1012.9	738.0	840.3	920.3	999.3
820.9	962.4	1095.9	1130.8	882.4	940.8	924.1	981.7	898.9	1031.0	1122.9	1187.9	898.8	1001.8	1070.7	1148.7
1129.8	1159.0	1214.7	1355.9	1120.1	1091.9	999.4	1051.1	1161.7	1227.4	1234.6	1363.3	1168.8	1189.4	1212.7	1386.6
1257.1	1237.4	1276.3	1469.4	1267.9	1102.4	1001.8	1050.4	1303.4	1259.5	1273.3	1337.2	1349.3	1305.3	1264.6	1458.0
1325.0	1231.3	1400.8	1492.7	1269.8	1132.8	1057.0	1025.4	1260.3	1294.3	1389.9	1448.3	1454.3	1402.8	1386.6	1502.4
1413.1	1466.0	1634.5	1647.7	1301.5	1228.8	1122.1	1161.9	1262.2	1369.0	1404.4	1443.1	1483.5	1461.5	1403.4	1460.7
חפוציות השימוש בתרופות אנטיביוטיות למיקוד (Antibiotics of concern)															
32.4%	31.5%	26.4%	29.6%	24.7%	25.2%	26.0%	27.7%	27.1%	28.9%	26.4%	27.3%	23.0%	25.0%	24.8%	29.3%
35.4%	35.2%	31.3%	35.7%	29.0%	28.5%	27.9%	33.5%	29.3%	32.6%	30.3%	30.3%	23.9%	25.6%	25.6%	34.5%
49.7%	51.6%	49.4%	53.8%	47.0%	44.9%	42.1%	46.5%	49.8%	49.6%	48.9%	50.0%	42.6%	43.7%	43.7%	51.7%
50.8%	54.8%	54.7%	58.1%	43.7%	45.5%	46.7%	52.8%	48.4%	52.6%	50.5%	55.2%	41.1%	44.0%	46.9%	54.1%
55.6%	57.8%	58.5%	61.6%	47.8%	50.8%	51.6%	55.5%	53.8%	56.7%	55.3%	58.8%	44.0%	48.1%	49.4%	55.5%
58.6%	60.2%	60.1%	63.7%	52.7%	54.4%	53.6%	58.4%	52.2%	56.2%	56.5%	58.2%	48.0%	50.3%	51.7%	56.4%
60.0%	62.2%	61.5%	64.5%	55.7%	55.1%	56.9%	58.6%	58.5%	59.4%	59.2%	59.9%	48.5%	50.6%	51.1%	54.8%
58.5%	59.8%	60.5%	62.7%	55.6%	56.6%	55.9%	55.6%	60.1%	60.2%	57.2%	58.1%	48.4%	50.5%	50.2%	52.8%
53.3%	58.7%	58.0%	63.5%	54.9%	54.7%	53.0%	58.0%	57.5%	58.4%	54.4%	56.1%	47.4%	50.1%	51.1%	51.3%
55.7%	57.1%	56.2%	64.2%	55.4%	55.9%	56.8%	56.1%	56.3%	63.1%	58.3%	57.8%	50.6%	50.8%	52.1%	53.5%
29.0%	29.5%	24.5%	27.5%	22.2%	23.0%	23.4%	25.5%	23.9%	26.2%	24.4%	24.9%	20.7%	22.8%	23.2%	27.1%
32.6%	31.6%	27.7%	32.2%	25.5%	25.6%	25.9%	29.2%	26.0%	30.4%	28.2%	28.6%	22.5%	24.3%	24.9%	33.5%
47.4%	48.2%	45.5%	49.1%	36.6%	36.5%	35.8%	40.0%	46.4%	45.1%	44.7%	45.6%	37.5%	39.4%	40.6%	48.0%
48.4%	51.7%	51.2%	54.5%	37.3%	38.5%	40.3%	44.0%	46.6%	49.4%	49.7%	49.9%	38.0%	41.8%	43.3%	49.3%
51.2%	55.4%	54.9%	58.9%	39.7%	43.1%	45.2%	49.2%	49.5%	54.0%	53.7%	54.0%	40.7%	44.7%	46.3%	51.1%
53.8%	56.6%	57.9%	60.6%	43.0%	46.0%	48.1%	51.8%	53.0%	55.3%	55.2%	55.9%	44.2%	46.1%	47.8%	51.8%
54.0%	57.0%	58.4%	60.9%	45.5%	47.9%	49.2%	50.0%	53.5%	56.3%	56.3%	56.4%	43.8%	45.8%	46.1%	48.9%
49.8%	52.4%	53.6%	55.5%	44.4%	45.8%	47.4%	48.6%	53.0%	53.8%	53.2%	52.5%	41.5%	43.1%	44.7%	46.9%
46.1%	52.1%	46.9%	56.6%	42.6%	45.0%	47.2%	47.4%	50.5%	49.9%	53.1%	50.9%	39.4%	42.3%	44.6%	47.0%
44.4%	49.7%	52.7%	54.9%	45.4%	48.2%	47.8%	50.2%	51.3%	52.2%	52.9%	55.5%	41.0%	43.6%	46.7%	48.9%
שיעור המבוטחים בעלי בדיקת RNA חיובית להפטיטיס C אשר קיבלו טיפול אנטי-ויראלי בתום השנים האחרונות															
NA*	72.2%	74.8%	66.7%	88.9%	80.3%	79.3%	78.2%	NA*	82.8%	73.4%	65.8%	74.6%	79.0%	73.3%	67.2%
NA*	73.8%	75.0%	NA*	76.7%	77.9%	76.9%	82.2%	NA*	68.1%	72.7%	74.3%	67.9%	79.7%	70.6%	67.0%
NA*	86.2%	82.1%	NA*	88.6%	87.5%	88.6%	82.8%	NA*	86.8%	79.1%	73.3%	78.7%	79.3%	78.1%	69.0%
NA*	80.5%	69.4%	NA*	74.3%	79.6%	75.2%	82.1%	NA*	59.4%	69.2%	75.8%	76.7%	75.9%	66.5%	76.3%

חא מסומן ב- * מייצג ערך מכנה נמוך מ-30 מבוטחים

בריאות הנפש																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																											
מאוחדת				מכבי				לאומית				כללית																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																															
8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	8+9+10	6+7	4+5	1+2+3	← SES	↓ מין																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																										
שיעור ההימצאות של מחלת נפש קשה (SMI) בקרב בני 18 ומעלה																	18-24	25-34	35-44	45-54	55-64	65+	18-24	25-34	35-44	45-54	55-64	65+	18-24	25-34	35-44	45-54	55-64	65+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																									
0.4%	0.6%	0.6%	0.6%	0.8%	1.2%	1.7%	1.8%	2.2%	2.3%	2.5%	3.5%	0.7%	1.1%	1.3%	0.7%			0.5%	0.8%	1.1%	1.2%	1.1%	1.5%	2.3%	2.7%	2.5%	2.9%	3.6%	3.8%	1.2%	1.7%	2.0%	1.6%			1.1%	1.3%	1.6%	1.4%	1.4%	2.0%	2.9%	3.5%	2.5%	3.1%	3.7%	4.0%	1.3%	2.0%	2.6%	2.2%			0.8%	1.2%	1.8%	1.5%	1.3%	2.0%	3.2%	3.5%	2.3%	3.0%	3.5%	3.8%	1.4%	2.3%	2.9%	2.4%			0.7%	0.8%	1.4%	1.5%	1.2%	1.7%	2.3%	2.9%	1.9%	2.1%	3.3%	4.3%	1.4%	2.4%	2.8%	2.7%			0.5%	0.6%	0.9%	0.8%	1.0%	1.3%	1.6%	1.9%	1.4%	1.5%	2.0%	2.3%	0.9%	1.3%	1.7%	1.6%			0.2%	0.2%	0.2%	0.1%	0.6%	0.9%	1.0%	0.5%	1.4%	1.3%	0.9%	0.8%	0.6%	0.6%	0.6%	0.3%			0.4%	0.5%	0.4%	0.4%	0.9%	1.1%	1.3%	1.2%	1.4%	1.9%	1.8%	1.5%	0.7%	0.9%	1.0%	0.8%			0.6%	0.7%	0.8%	0.8%	1.1%	1.5%	2.0%	2.0%	1.6%	1.7%	2.5%	2.5%	0.8%	1.3%	1.5%	1.3%			0.7%	0.8%	1.1%	1.1%	1.1%	1.7%	2.3%	2.5%	1.7%	2.3%	2.8%	2.6%	1.0%	1.7%	2.1%	1.7%			0.7%	0.8%	1.3%	1.0%	1.3%	1.7%	2.4%	2.8%	1.9%	2.1%	2.8%	3.1%	1.3%	1.9%	2.4%	2.0%			0.8%	0.8%	0.9%	0.9%	1.3%	1.6%	1.9%	2.1%	1.9%	1.9%	2.1%	2.7%	1.1%	1.4%	1.7%	1.6%			שיעור ההימצאות של סוכרת בקרב חולי SMI בני 18 ומעלה																	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	1.7%	2.5%	1.3%	1.8%	2.0%	2.2%	2.6%	3.9%	0.0%	1.1%	1.3%	1.2%	2.2%	1.7%	1.9%	2.7%			2.3%	4.6%	6.5%	7.0%	5.5%	5.8%	7.0%	7.8%	0.0%	4.5%	6.3%	6.2%	6.9%	6.6%	8.7%	9.4%			17.4%	16.0%	17.0%	18.8%	12.3%	14.1%	17.7%	19.0%	7.7%	10.6%	14.9%	17.4%	14.0%	16.2%	16.7%	19.1%			11.8%	29.3%	26.4%	34.4%	20.3%	26.0%	27.2%	30.5%	30.6%	26.8%	25.7%	23.4%	24.0%	27.3%	28.7%	31.2%			39.6%	33.3%	27.3%	36.5%	28.4%	42.0%	33.3%	34.8%	43.2%	41.2%	35.5%	36.2%	36.4%	33.2%	35.0%	33.2%			0.0%	5.5%	3.0%	2.1%	3.2%	2.2%	3.8%	2.3%	1.4%	1.8%	1.4%	1.6%	2.0%	2.6%	4.2%	4.0%			11.8%	6.0%	8.5%	9.1%	5.1%	7.7%	9.5%	9.7%	5.0%	5.0%	3.4%	9.4%	4.5%	6.6%	7.3%	9.8%			8.5%	9.6%	15.3%	18.5%	10.0%	13.5%	17.2%	20.6%	11.1%	10.5%	12.6%	16.7%	10.6%	15.7%	18.2%	21.1%			28.1%	34.5%	27.4%	32.3%	17.9%	21.0%	27.0%	29.2%	22.7%	21.9%	30.0%	24.2%	20.8%	28.3%	29.3%	33.5%			21.6%	30.2%	33.7%	36.8%	25.7%	32.7%	33.0%	34.2%	27.8%	25.5%	32.5%	33.5%	28.9%	34.4%	36.9%	39.9%			שיעור בעלי יתום HbA1c מאוזנת בחולי סוכרת ו SMI בני 18-84																	18-44	45-64	65+	18-44	45-64	65+	18-44	45-64	65+	NA*	NA*	NA*	NA*	83.7%	67.7%	74.3%	70.7%	NA*	NA*	59.4%	NA*	76.7%	73.5%	72.4%	70.6%	18-44	NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+
0.5%	0.8%	1.1%	1.2%	1.1%	1.5%	2.3%	2.7%	2.5%	2.9%	3.6%	3.8%	1.2%	1.7%	2.0%	1.6%			1.1%	1.3%	1.6%	1.4%	1.4%	2.0%	2.9%	3.5%	2.5%	3.1%	3.7%	4.0%	1.3%	2.0%	2.6%	2.2%			0.8%	1.2%	1.8%	1.5%	1.3%	2.0%	3.2%	3.5%	2.3%	3.0%	3.5%	3.8%	1.4%	2.3%	2.9%	2.4%			0.7%	0.8%	1.4%	1.5%	1.2%	1.7%	2.3%	2.9%	1.9%	2.1%	3.3%	4.3%	1.4%	2.4%	2.8%	2.7%			0.5%	0.6%	0.9%	0.8%	1.0%	1.3%	1.6%	1.9%	1.4%	1.5%	2.0%	2.3%	0.9%	1.3%	1.7%	1.6%			0.2%	0.2%	0.2%	0.1%	0.6%	0.9%	1.0%	0.5%	1.4%	1.3%	0.9%	0.8%	0.6%	0.6%	0.6%	0.3%			0.4%	0.5%	0.4%	0.4%	0.9%	1.1%	1.3%	1.2%	1.4%	1.9%	1.8%	1.5%	0.7%	0.9%	1.0%	0.8%			0.6%	0.7%	0.8%	0.8%	1.1%	1.5%	2.0%	2.0%	1.6%	1.7%	2.5%	2.5%	0.8%	1.3%	1.5%	1.3%			0.7%	0.8%	1.1%	1.1%	1.1%	1.7%	2.3%	2.5%	1.7%	2.3%	2.8%	2.6%	1.0%	1.7%	2.1%	1.7%			0.7%	0.8%	1.3%	1.0%	1.3%	1.7%	2.4%	2.8%	1.9%	2.1%	2.8%	3.1%	1.3%	1.9%	2.4%	2.0%			0.8%	0.8%	0.9%	0.9%	1.3%	1.6%	1.9%	2.1%	1.9%	1.9%	2.1%	2.7%	1.1%	1.4%	1.7%	1.6%			שיעור ההימצאות של סוכרת בקרב חולי SMI בני 18 ומעלה																	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	1.7%	2.5%	1.3%	1.8%	2.0%	2.2%	2.6%	3.9%	0.0%	1.1%	1.3%	1.2%	2.2%	1.7%	1.9%	2.7%			2.3%	4.6%	6.5%	7.0%	5.5%	5.8%	7.0%	7.8%	0.0%	4.5%	6.3%	6.2%	6.9%	6.6%	8.7%	9.4%			17.4%	16.0%	17.0%	18.8%	12.3%	14.1%	17.7%	19.0%	7.7%	10.6%	14.9%	17.4%	14.0%	16.2%	16.7%	19.1%			11.8%	29.3%	26.4%	34.4%	20.3%	26.0%	27.2%	30.5%	30.6%	26.8%	25.7%	23.4%	24.0%	27.3%	28.7%	31.2%			39.6%	33.3%	27.3%	36.5%	28.4%	42.0%	33.3%	34.8%	43.2%	41.2%	35.5%	36.2%	36.4%	33.2%	35.0%	33.2%			0.0%	5.5%	3.0%	2.1%	3.2%	2.2%	3.8%	2.3%	1.4%	1.8%	1.4%	1.6%	2.0%	2.6%	4.2%	4.0%			11.8%	6.0%	8.5%	9.1%	5.1%	7.7%	9.5%	9.7%	5.0%	5.0%	3.4%	9.4%	4.5%	6.6%	7.3%	9.8%			8.5%	9.6%	15.3%	18.5%	10.0%	13.5%	17.2%	20.6%	11.1%	10.5%	12.6%	16.7%	10.6%	15.7%	18.2%	21.1%			28.1%	34.5%	27.4%	32.3%	17.9%	21.0%	27.0%	29.2%	22.7%	21.9%	30.0%	24.2%	20.8%	28.3%	29.3%	33.5%			21.6%	30.2%	33.7%	36.8%	25.7%	32.7%	33.0%	34.2%	27.8%	25.5%	32.5%	33.5%	28.9%	34.4%	36.9%	39.9%			שיעור בעלי יתום HbA1c מאוזנת בחולי סוכרת ו SMI בני 18-84																	18-44	45-64	65+	18-44	45-64	65+	18-44	45-64	65+	NA*	NA*	NA*	NA*	83.7%	67.7%	74.3%	70.7%	NA*	NA*	59.4%	NA*	76.7%	73.5%	72.4%	70.6%	18-44	NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																		
1.1%	1.3%	1.6%	1.4%	1.4%	2.0%	2.9%	3.5%	2.5%	3.1%	3.7%	4.0%	1.3%	2.0%	2.6%	2.2%			0.8%	1.2%	1.8%	1.5%	1.3%	2.0%	3.2%	3.5%	2.3%	3.0%	3.5%	3.8%	1.4%	2.3%	2.9%	2.4%			0.7%	0.8%	1.4%	1.5%	1.2%	1.7%	2.3%	2.9%	1.9%	2.1%	3.3%	4.3%	1.4%	2.4%	2.8%	2.7%			0.5%	0.6%	0.9%	0.8%	1.0%	1.3%	1.6%	1.9%	1.4%	1.5%	2.0%	2.3%	0.9%	1.3%	1.7%	1.6%			0.2%	0.2%	0.2%	0.1%	0.6%	0.9%	1.0%	0.5%	1.4%	1.3%	0.9%	0.8%	0.6%	0.6%	0.6%	0.3%			0.4%	0.5%	0.4%	0.4%	0.9%	1.1%	1.3%	1.2%	1.4%	1.9%	1.8%	1.5%	0.7%	0.9%	1.0%	0.8%			0.6%	0.7%	0.8%	0.8%	1.1%	1.5%	2.0%	2.0%	1.6%	1.7%	2.5%	2.5%	0.8%	1.3%	1.5%	1.3%			0.7%	0.8%	1.1%	1.1%	1.1%	1.7%	2.3%	2.5%	1.7%	2.3%	2.8%	2.6%	1.0%	1.7%	2.1%	1.7%			0.7%	0.8%	1.3%	1.0%	1.3%	1.7%	2.4%	2.8%	1.9%	2.1%	2.8%	3.1%	1.3%	1.9%	2.4%	2.0%			0.8%	0.8%	0.9%	0.9%	1.3%	1.6%	1.9%	2.1%	1.9%	1.9%	2.1%	2.7%	1.1%	1.4%	1.7%	1.6%			שיעור ההימצאות של סוכרת בקרב חולי SMI בני 18 ומעלה																	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	1.7%	2.5%	1.3%	1.8%	2.0%	2.2%	2.6%	3.9%	0.0%	1.1%	1.3%	1.2%	2.2%	1.7%	1.9%	2.7%			2.3%	4.6%	6.5%	7.0%	5.5%	5.8%	7.0%	7.8%	0.0%	4.5%	6.3%	6.2%	6.9%	6.6%	8.7%	9.4%			17.4%	16.0%	17.0%	18.8%	12.3%	14.1%	17.7%	19.0%	7.7%	10.6%	14.9%	17.4%	14.0%	16.2%	16.7%	19.1%			11.8%	29.3%	26.4%	34.4%	20.3%	26.0%	27.2%	30.5%	30.6%	26.8%	25.7%	23.4%	24.0%	27.3%	28.7%	31.2%			39.6%	33.3%	27.3%	36.5%	28.4%	42.0%	33.3%	34.8%	43.2%	41.2%	35.5%	36.2%	36.4%	33.2%	35.0%	33.2%			0.0%	5.5%	3.0%	2.1%	3.2%	2.2%	3.8%	2.3%	1.4%	1.8%	1.4%	1.6%	2.0%	2.6%	4.2%	4.0%			11.8%	6.0%	8.5%	9.1%	5.1%	7.7%	9.5%	9.7%	5.0%	5.0%	3.4%	9.4%	4.5%	6.6%	7.3%	9.8%			8.5%	9.6%	15.3%	18.5%	10.0%	13.5%	17.2%	20.6%	11.1%	10.5%	12.6%	16.7%	10.6%	15.7%	18.2%	21.1%			28.1%	34.5%	27.4%	32.3%	17.9%	21.0%	27.0%	29.2%	22.7%	21.9%	30.0%	24.2%	20.8%	28.3%	29.3%	33.5%			21.6%	30.2%	33.7%	36.8%	25.7%	32.7%	33.0%	34.2%	27.8%	25.5%	32.5%	33.5%	28.9%	34.4%	36.9%	39.9%			שיעור בעלי יתום HbA1c מאוזנת בחולי סוכרת ו SMI בני 18-84																	18-44	45-64	65+	18-44	45-64	65+	18-44	45-64	65+	NA*	NA*	NA*	NA*	83.7%	67.7%	74.3%	70.7%	NA*	NA*	59.4%	NA*	76.7%	73.5%	72.4%	70.6%	18-44	NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																				
0.8%	1.2%	1.8%	1.5%	1.3%	2.0%	3.2%	3.5%	2.3%	3.0%	3.5%	3.8%	1.4%	2.3%	2.9%	2.4%			0.7%	0.8%	1.4%	1.5%	1.2%	1.7%	2.3%	2.9%	1.9%	2.1%	3.3%	4.3%	1.4%	2.4%	2.8%	2.7%			0.5%	0.6%	0.9%	0.8%	1.0%	1.3%	1.6%	1.9%	1.4%	1.5%	2.0%	2.3%	0.9%	1.3%	1.7%	1.6%			0.2%	0.2%	0.2%	0.1%	0.6%	0.9%	1.0%	0.5%	1.4%	1.3%	0.9%	0.8%	0.6%	0.6%	0.6%	0.3%			0.4%	0.5%	0.4%	0.4%	0.9%	1.1%	1.3%	1.2%	1.4%	1.9%	1.8%	1.5%	0.7%	0.9%	1.0%	0.8%			0.6%	0.7%	0.8%	0.8%	1.1%	1.5%	2.0%	2.0%	1.6%	1.7%	2.5%	2.5%	0.8%	1.3%	1.5%	1.3%			0.7%	0.8%	1.1%	1.1%	1.1%	1.7%	2.3%	2.5%	1.7%	2.3%	2.8%	2.6%	1.0%	1.7%	2.1%	1.7%			0.7%	0.8%	1.3%	1.0%	1.3%	1.7%	2.4%	2.8%	1.9%	2.1%	2.8%	3.1%	1.3%	1.9%	2.4%	2.0%			0.8%	0.8%	0.9%	0.9%	1.3%	1.6%	1.9%	2.1%	1.9%	1.9%	2.1%	2.7%	1.1%	1.4%	1.7%	1.6%			שיעור ההימצאות של סוכרת בקרב חולי SMI בני 18 ומעלה																	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	1.7%	2.5%	1.3%	1.8%	2.0%	2.2%	2.6%	3.9%	0.0%	1.1%	1.3%	1.2%	2.2%	1.7%	1.9%	2.7%			2.3%	4.6%	6.5%	7.0%	5.5%	5.8%	7.0%	7.8%	0.0%	4.5%	6.3%	6.2%	6.9%	6.6%	8.7%	9.4%			17.4%	16.0%	17.0%	18.8%	12.3%	14.1%	17.7%	19.0%	7.7%	10.6%	14.9%	17.4%	14.0%	16.2%	16.7%	19.1%			11.8%	29.3%	26.4%	34.4%	20.3%	26.0%	27.2%	30.5%	30.6%	26.8%	25.7%	23.4%	24.0%	27.3%	28.7%	31.2%			39.6%	33.3%	27.3%	36.5%	28.4%	42.0%	33.3%	34.8%	43.2%	41.2%	35.5%	36.2%	36.4%	33.2%	35.0%	33.2%			0.0%	5.5%	3.0%	2.1%	3.2%	2.2%	3.8%	2.3%	1.4%	1.8%	1.4%	1.6%	2.0%	2.6%	4.2%	4.0%			11.8%	6.0%	8.5%	9.1%	5.1%	7.7%	9.5%	9.7%	5.0%	5.0%	3.4%	9.4%	4.5%	6.6%	7.3%	9.8%			8.5%	9.6%	15.3%	18.5%	10.0%	13.5%	17.2%	20.6%	11.1%	10.5%	12.6%	16.7%	10.6%	15.7%	18.2%	21.1%			28.1%	34.5%	27.4%	32.3%	17.9%	21.0%	27.0%	29.2%	22.7%	21.9%	30.0%	24.2%	20.8%	28.3%	29.3%	33.5%			21.6%	30.2%	33.7%	36.8%	25.7%	32.7%	33.0%	34.2%	27.8%	25.5%	32.5%	33.5%	28.9%	34.4%	36.9%	39.9%			שיעור בעלי יתום HbA1c מאוזנת בחולי סוכרת ו SMI בני 18-84																	18-44	45-64	65+	18-44	45-64	65+	18-44	45-64	65+	NA*	NA*	NA*	NA*	83.7%	67.7%	74.3%	70.7%	NA*	NA*	59.4%	NA*	76.7%	73.5%	72.4%	70.6%	18-44	NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																						
0.7%	0.8%	1.4%	1.5%	1.2%	1.7%	2.3%	2.9%	1.9%	2.1%	3.3%	4.3%	1.4%	2.4%	2.8%	2.7%			0.5%	0.6%	0.9%	0.8%	1.0%	1.3%	1.6%	1.9%	1.4%	1.5%	2.0%	2.3%	0.9%	1.3%	1.7%	1.6%			0.2%	0.2%	0.2%	0.1%	0.6%	0.9%	1.0%	0.5%	1.4%	1.3%	0.9%	0.8%	0.6%	0.6%	0.6%	0.3%			0.4%	0.5%	0.4%	0.4%	0.9%	1.1%	1.3%	1.2%	1.4%	1.9%	1.8%	1.5%	0.7%	0.9%	1.0%	0.8%			0.6%	0.7%	0.8%	0.8%	1.1%	1.5%	2.0%	2.0%	1.6%	1.7%	2.5%	2.5%	0.8%	1.3%	1.5%	1.3%			0.7%	0.8%	1.1%	1.1%	1.1%	1.7%	2.3%	2.5%	1.7%	2.3%	2.8%	2.6%	1.0%	1.7%	2.1%	1.7%			0.7%	0.8%	1.3%	1.0%	1.3%	1.7%	2.4%	2.8%	1.9%	2.1%	2.8%	3.1%	1.3%	1.9%	2.4%	2.0%			0.8%	0.8%	0.9%	0.9%	1.3%	1.6%	1.9%	2.1%	1.9%	1.9%	2.1%	2.7%	1.1%	1.4%	1.7%	1.6%			שיעור ההימצאות של סוכרת בקרב חולי SMI בני 18 ומעלה																	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	1.7%	2.5%	1.3%	1.8%	2.0%	2.2%	2.6%	3.9%	0.0%	1.1%	1.3%	1.2%	2.2%	1.7%	1.9%	2.7%			2.3%	4.6%	6.5%	7.0%	5.5%	5.8%	7.0%	7.8%	0.0%	4.5%	6.3%	6.2%	6.9%	6.6%	8.7%	9.4%			17.4%	16.0%	17.0%	18.8%	12.3%	14.1%	17.7%	19.0%	7.7%	10.6%	14.9%	17.4%	14.0%	16.2%	16.7%	19.1%			11.8%	29.3%	26.4%	34.4%	20.3%	26.0%	27.2%	30.5%	30.6%	26.8%	25.7%	23.4%	24.0%	27.3%	28.7%	31.2%			39.6%	33.3%	27.3%	36.5%	28.4%	42.0%	33.3%	34.8%	43.2%	41.2%	35.5%	36.2%	36.4%	33.2%	35.0%	33.2%			0.0%	5.5%	3.0%	2.1%	3.2%	2.2%	3.8%	2.3%	1.4%	1.8%	1.4%	1.6%	2.0%	2.6%	4.2%	4.0%			11.8%	6.0%	8.5%	9.1%	5.1%	7.7%	9.5%	9.7%	5.0%	5.0%	3.4%	9.4%	4.5%	6.6%	7.3%	9.8%			8.5%	9.6%	15.3%	18.5%	10.0%	13.5%	17.2%	20.6%	11.1%	10.5%	12.6%	16.7%	10.6%	15.7%	18.2%	21.1%			28.1%	34.5%	27.4%	32.3%	17.9%	21.0%	27.0%	29.2%	22.7%	21.9%	30.0%	24.2%	20.8%	28.3%	29.3%	33.5%			21.6%	30.2%	33.7%	36.8%	25.7%	32.7%	33.0%	34.2%	27.8%	25.5%	32.5%	33.5%	28.9%	34.4%	36.9%	39.9%			שיעור בעלי יתום HbA1c מאוזנת בחולי סוכרת ו SMI בני 18-84																	18-44	45-64	65+	18-44	45-64	65+	18-44	45-64	65+	NA*	NA*	NA*	NA*	83.7%	67.7%	74.3%	70.7%	NA*	NA*	59.4%	NA*	76.7%	73.5%	72.4%	70.6%	18-44	NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																								
0.5%	0.6%	0.9%	0.8%	1.0%	1.3%	1.6%	1.9%	1.4%	1.5%	2.0%	2.3%	0.9%	1.3%	1.7%	1.6%			0.2%	0.2%	0.2%	0.1%	0.6%	0.9%	1.0%	0.5%	1.4%	1.3%	0.9%	0.8%	0.6%	0.6%	0.6%	0.3%			0.4%	0.5%	0.4%	0.4%	0.9%	1.1%	1.3%	1.2%	1.4%	1.9%	1.8%	1.5%	0.7%	0.9%	1.0%	0.8%			0.6%	0.7%	0.8%	0.8%	1.1%	1.5%	2.0%	2.0%	1.6%	1.7%	2.5%	2.5%	0.8%	1.3%	1.5%	1.3%			0.7%	0.8%	1.1%	1.1%	1.1%	1.7%	2.3%	2.5%	1.7%	2.3%	2.8%	2.6%	1.0%	1.7%	2.1%	1.7%			0.7%	0.8%	1.3%	1.0%	1.3%	1.7%	2.4%	2.8%	1.9%	2.1%	2.8%	3.1%	1.3%	1.9%	2.4%	2.0%			0.8%	0.8%	0.9%	0.9%	1.3%	1.6%	1.9%	2.1%	1.9%	1.9%	2.1%	2.7%	1.1%	1.4%	1.7%	1.6%			שיעור ההימצאות של סוכרת בקרב חולי SMI בני 18 ומעלה																	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	1.7%	2.5%	1.3%	1.8%	2.0%	2.2%	2.6%	3.9%	0.0%	1.1%	1.3%	1.2%	2.2%	1.7%	1.9%	2.7%			2.3%	4.6%	6.5%	7.0%	5.5%	5.8%	7.0%	7.8%	0.0%	4.5%	6.3%	6.2%	6.9%	6.6%	8.7%	9.4%			17.4%	16.0%	17.0%	18.8%	12.3%	14.1%	17.7%	19.0%	7.7%	10.6%	14.9%	17.4%	14.0%	16.2%	16.7%	19.1%			11.8%	29.3%	26.4%	34.4%	20.3%	26.0%	27.2%	30.5%	30.6%	26.8%	25.7%	23.4%	24.0%	27.3%	28.7%	31.2%			39.6%	33.3%	27.3%	36.5%	28.4%	42.0%	33.3%	34.8%	43.2%	41.2%	35.5%	36.2%	36.4%	33.2%	35.0%	33.2%			0.0%	5.5%	3.0%	2.1%	3.2%	2.2%	3.8%	2.3%	1.4%	1.8%	1.4%	1.6%	2.0%	2.6%	4.2%	4.0%			11.8%	6.0%	8.5%	9.1%	5.1%	7.7%	9.5%	9.7%	5.0%	5.0%	3.4%	9.4%	4.5%	6.6%	7.3%	9.8%			8.5%	9.6%	15.3%	18.5%	10.0%	13.5%	17.2%	20.6%	11.1%	10.5%	12.6%	16.7%	10.6%	15.7%	18.2%	21.1%			28.1%	34.5%	27.4%	32.3%	17.9%	21.0%	27.0%	29.2%	22.7%	21.9%	30.0%	24.2%	20.8%	28.3%	29.3%	33.5%			21.6%	30.2%	33.7%	36.8%	25.7%	32.7%	33.0%	34.2%	27.8%	25.5%	32.5%	33.5%	28.9%	34.4%	36.9%	39.9%			שיעור בעלי יתום HbA1c מאוזנת בחולי סוכרת ו SMI בני 18-84																	18-44	45-64	65+	18-44	45-64	65+	18-44	45-64	65+	NA*	NA*	NA*	NA*	83.7%	67.7%	74.3%	70.7%	NA*	NA*	59.4%	NA*	76.7%	73.5%	72.4%	70.6%	18-44	NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																										
0.2%	0.2%	0.2%	0.1%	0.6%	0.9%	1.0%	0.5%	1.4%	1.3%	0.9%	0.8%	0.6%	0.6%	0.6%	0.3%			0.4%	0.5%	0.4%	0.4%	0.9%	1.1%	1.3%	1.2%	1.4%	1.9%	1.8%	1.5%	0.7%	0.9%	1.0%	0.8%			0.6%	0.7%	0.8%	0.8%	1.1%	1.5%	2.0%	2.0%	1.6%	1.7%	2.5%	2.5%	0.8%	1.3%	1.5%	1.3%			0.7%	0.8%	1.1%	1.1%	1.1%	1.7%	2.3%	2.5%	1.7%	2.3%	2.8%	2.6%	1.0%	1.7%	2.1%	1.7%			0.7%	0.8%	1.3%	1.0%	1.3%	1.7%	2.4%	2.8%	1.9%	2.1%	2.8%	3.1%	1.3%	1.9%	2.4%	2.0%			0.8%	0.8%	0.9%	0.9%	1.3%	1.6%	1.9%	2.1%	1.9%	1.9%	2.1%	2.7%	1.1%	1.4%	1.7%	1.6%			שיעור ההימצאות של סוכרת בקרב חולי SMI בני 18 ומעלה																	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	1.7%	2.5%	1.3%	1.8%	2.0%	2.2%	2.6%	3.9%	0.0%	1.1%	1.3%	1.2%	2.2%	1.7%	1.9%	2.7%			2.3%	4.6%	6.5%	7.0%	5.5%	5.8%	7.0%	7.8%	0.0%	4.5%	6.3%	6.2%	6.9%	6.6%	8.7%	9.4%			17.4%	16.0%	17.0%	18.8%	12.3%	14.1%	17.7%	19.0%	7.7%	10.6%	14.9%	17.4%	14.0%	16.2%	16.7%	19.1%			11.8%	29.3%	26.4%	34.4%	20.3%	26.0%	27.2%	30.5%	30.6%	26.8%	25.7%	23.4%	24.0%	27.3%	28.7%	31.2%			39.6%	33.3%	27.3%	36.5%	28.4%	42.0%	33.3%	34.8%	43.2%	41.2%	35.5%	36.2%	36.4%	33.2%	35.0%	33.2%			0.0%	5.5%	3.0%	2.1%	3.2%	2.2%	3.8%	2.3%	1.4%	1.8%	1.4%	1.6%	2.0%	2.6%	4.2%	4.0%			11.8%	6.0%	8.5%	9.1%	5.1%	7.7%	9.5%	9.7%	5.0%	5.0%	3.4%	9.4%	4.5%	6.6%	7.3%	9.8%			8.5%	9.6%	15.3%	18.5%	10.0%	13.5%	17.2%	20.6%	11.1%	10.5%	12.6%	16.7%	10.6%	15.7%	18.2%	21.1%			28.1%	34.5%	27.4%	32.3%	17.9%	21.0%	27.0%	29.2%	22.7%	21.9%	30.0%	24.2%	20.8%	28.3%	29.3%	33.5%			21.6%	30.2%	33.7%	36.8%	25.7%	32.7%	33.0%	34.2%	27.8%	25.5%	32.5%	33.5%	28.9%	34.4%	36.9%	39.9%			שיעור בעלי יתום HbA1c מאוזנת בחולי סוכרת ו SMI בני 18-84																	18-44	45-64	65+	18-44	45-64	65+	18-44	45-64	65+	NA*	NA*	NA*	NA*	83.7%	67.7%	74.3%	70.7%	NA*	NA*	59.4%	NA*	76.7%	73.5%	72.4%	70.6%	18-44	NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																												
0.4%	0.5%	0.4%	0.4%	0.9%	1.1%	1.3%	1.2%	1.4%	1.9%	1.8%	1.5%	0.7%	0.9%	1.0%	0.8%			0.6%	0.7%	0.8%	0.8%	1.1%	1.5%	2.0%	2.0%	1.6%	1.7%	2.5%	2.5%	0.8%	1.3%	1.5%	1.3%			0.7%	0.8%	1.1%	1.1%	1.1%	1.7%	2.3%	2.5%	1.7%	2.3%	2.8%	2.6%	1.0%	1.7%	2.1%	1.7%			0.7%	0.8%	1.3%	1.0%	1.3%	1.7%	2.4%	2.8%	1.9%	2.1%	2.8%	3.1%	1.3%	1.9%	2.4%	2.0%			0.8%	0.8%	0.9%	0.9%	1.3%	1.6%	1.9%	2.1%	1.9%	1.9%	2.1%	2.7%	1.1%	1.4%	1.7%	1.6%			שיעור ההימצאות של סוכרת בקרב חולי SMI בני 18 ומעלה																	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	1.7%	2.5%	1.3%	1.8%	2.0%	2.2%	2.6%	3.9%	0.0%	1.1%	1.3%	1.2%	2.2%	1.7%	1.9%	2.7%			2.3%	4.6%	6.5%	7.0%	5.5%	5.8%	7.0%	7.8%	0.0%	4.5%	6.3%	6.2%	6.9%	6.6%	8.7%	9.4%			17.4%	16.0%	17.0%	18.8%	12.3%	14.1%	17.7%	19.0%	7.7%	10.6%	14.9%	17.4%	14.0%	16.2%	16.7%	19.1%			11.8%	29.3%	26.4%	34.4%	20.3%	26.0%	27.2%	30.5%	30.6%	26.8%	25.7%	23.4%	24.0%	27.3%	28.7%	31.2%			39.6%	33.3%	27.3%	36.5%	28.4%	42.0%	33.3%	34.8%	43.2%	41.2%	35.5%	36.2%	36.4%	33.2%	35.0%	33.2%			0.0%	5.5%	3.0%	2.1%	3.2%	2.2%	3.8%	2.3%	1.4%	1.8%	1.4%	1.6%	2.0%	2.6%	4.2%	4.0%			11.8%	6.0%	8.5%	9.1%	5.1%	7.7%	9.5%	9.7%	5.0%	5.0%	3.4%	9.4%	4.5%	6.6%	7.3%	9.8%			8.5%	9.6%	15.3%	18.5%	10.0%	13.5%	17.2%	20.6%	11.1%	10.5%	12.6%	16.7%	10.6%	15.7%	18.2%	21.1%			28.1%	34.5%	27.4%	32.3%	17.9%	21.0%	27.0%	29.2%	22.7%	21.9%	30.0%	24.2%	20.8%	28.3%	29.3%	33.5%			21.6%	30.2%	33.7%	36.8%	25.7%	32.7%	33.0%	34.2%	27.8%	25.5%	32.5%	33.5%	28.9%	34.4%	36.9%	39.9%			שיעור בעלי יתום HbA1c מאוזנת בחולי סוכרת ו SMI בני 18-84																	18-44	45-64	65+	18-44	45-64	65+	18-44	45-64	65+	NA*	NA*	NA*	NA*	83.7%	67.7%	74.3%	70.7%	NA*	NA*	59.4%	NA*	76.7%	73.5%	72.4%	70.6%	18-44	NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																														
0.6%	0.7%	0.8%	0.8%	1.1%	1.5%	2.0%	2.0%	1.6%	1.7%	2.5%	2.5%	0.8%	1.3%	1.5%	1.3%			0.7%	0.8%	1.1%	1.1%	1.1%	1.7%	2.3%	2.5%	1.7%	2.3%	2.8%	2.6%	1.0%	1.7%	2.1%	1.7%			0.7%	0.8%	1.3%	1.0%	1.3%	1.7%	2.4%	2.8%	1.9%	2.1%	2.8%	3.1%	1.3%	1.9%	2.4%	2.0%			0.8%	0.8%	0.9%	0.9%	1.3%	1.6%	1.9%	2.1%	1.9%	1.9%	2.1%	2.7%	1.1%	1.4%	1.7%	1.6%			שיעור ההימצאות של סוכרת בקרב חולי SMI בני 18 ומעלה																	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	1.7%	2.5%	1.3%	1.8%	2.0%	2.2%	2.6%	3.9%	0.0%	1.1%	1.3%	1.2%	2.2%	1.7%	1.9%	2.7%			2.3%	4.6%	6.5%	7.0%	5.5%	5.8%	7.0%	7.8%	0.0%	4.5%	6.3%	6.2%	6.9%	6.6%	8.7%	9.4%			17.4%	16.0%	17.0%	18.8%	12.3%	14.1%	17.7%	19.0%	7.7%	10.6%	14.9%	17.4%	14.0%	16.2%	16.7%	19.1%			11.8%	29.3%	26.4%	34.4%	20.3%	26.0%	27.2%	30.5%	30.6%	26.8%	25.7%	23.4%	24.0%	27.3%	28.7%	31.2%			39.6%	33.3%	27.3%	36.5%	28.4%	42.0%	33.3%	34.8%	43.2%	41.2%	35.5%	36.2%	36.4%	33.2%	35.0%	33.2%			0.0%	5.5%	3.0%	2.1%	3.2%	2.2%	3.8%	2.3%	1.4%	1.8%	1.4%	1.6%	2.0%	2.6%	4.2%	4.0%			11.8%	6.0%	8.5%	9.1%	5.1%	7.7%	9.5%	9.7%	5.0%	5.0%	3.4%	9.4%	4.5%	6.6%	7.3%	9.8%			8.5%	9.6%	15.3%	18.5%	10.0%	13.5%	17.2%	20.6%	11.1%	10.5%	12.6%	16.7%	10.6%	15.7%	18.2%	21.1%			28.1%	34.5%	27.4%	32.3%	17.9%	21.0%	27.0%	29.2%	22.7%	21.9%	30.0%	24.2%	20.8%	28.3%	29.3%	33.5%			21.6%	30.2%	33.7%	36.8%	25.7%	32.7%	33.0%	34.2%	27.8%	25.5%	32.5%	33.5%	28.9%	34.4%	36.9%	39.9%			שיעור בעלי יתום HbA1c מאוזנת בחולי סוכרת ו SMI בני 18-84																	18-44	45-64	65+	18-44	45-64	65+	18-44	45-64	65+	NA*	NA*	NA*	NA*	83.7%	67.7%	74.3%	70.7%	NA*	NA*	59.4%	NA*	76.7%	73.5%	72.4%	70.6%	18-44	NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																
0.7%	0.8%	1.1%	1.1%	1.1%	1.7%	2.3%	2.5%	1.7%	2.3%	2.8%	2.6%	1.0%	1.7%	2.1%	1.7%			0.7%	0.8%	1.3%	1.0%	1.3%	1.7%	2.4%	2.8%	1.9%	2.1%	2.8%	3.1%	1.3%	1.9%	2.4%	2.0%			0.8%	0.8%	0.9%	0.9%	1.3%	1.6%	1.9%	2.1%	1.9%	1.9%	2.1%	2.7%	1.1%	1.4%	1.7%	1.6%			שיעור ההימצאות של סוכרת בקרב חולי SMI בני 18 ומעלה																	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	1.7%	2.5%	1.3%	1.8%	2.0%	2.2%	2.6%	3.9%	0.0%	1.1%	1.3%	1.2%	2.2%	1.7%	1.9%	2.7%			2.3%	4.6%	6.5%	7.0%	5.5%	5.8%	7.0%	7.8%	0.0%	4.5%	6.3%	6.2%	6.9%	6.6%	8.7%	9.4%			17.4%	16.0%	17.0%	18.8%	12.3%	14.1%	17.7%	19.0%	7.7%	10.6%	14.9%	17.4%	14.0%	16.2%	16.7%	19.1%			11.8%	29.3%	26.4%	34.4%	20.3%	26.0%	27.2%	30.5%	30.6%	26.8%	25.7%	23.4%	24.0%	27.3%	28.7%	31.2%			39.6%	33.3%	27.3%	36.5%	28.4%	42.0%	33.3%	34.8%	43.2%	41.2%	35.5%	36.2%	36.4%	33.2%	35.0%	33.2%			0.0%	5.5%	3.0%	2.1%	3.2%	2.2%	3.8%	2.3%	1.4%	1.8%	1.4%	1.6%	2.0%	2.6%	4.2%	4.0%			11.8%	6.0%	8.5%	9.1%	5.1%	7.7%	9.5%	9.7%	5.0%	5.0%	3.4%	9.4%	4.5%	6.6%	7.3%	9.8%			8.5%	9.6%	15.3%	18.5%	10.0%	13.5%	17.2%	20.6%	11.1%	10.5%	12.6%	16.7%	10.6%	15.7%	18.2%	21.1%			28.1%	34.5%	27.4%	32.3%	17.9%	21.0%	27.0%	29.2%	22.7%	21.9%	30.0%	24.2%	20.8%	28.3%	29.3%	33.5%			21.6%	30.2%	33.7%	36.8%	25.7%	32.7%	33.0%	34.2%	27.8%	25.5%	32.5%	33.5%	28.9%	34.4%	36.9%	39.9%			שיעור בעלי יתום HbA1c מאוזנת בחולי סוכרת ו SMI בני 18-84																	18-44	45-64	65+	18-44	45-64	65+	18-44	45-64	65+	NA*	NA*	NA*	NA*	83.7%	67.7%	74.3%	70.7%	NA*	NA*	59.4%	NA*	76.7%	73.5%	72.4%	70.6%	18-44	NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																		
0.7%	0.8%	1.3%	1.0%	1.3%	1.7%	2.4%	2.8%	1.9%	2.1%	2.8%	3.1%	1.3%	1.9%	2.4%	2.0%			0.8%	0.8%	0.9%	0.9%	1.3%	1.6%	1.9%	2.1%	1.9%	1.9%	2.1%	2.7%	1.1%	1.4%	1.7%	1.6%			שיעור ההימצאות של סוכרת בקרב חולי SMI בני 18 ומעלה																	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	1.7%	2.5%	1.3%	1.8%	2.0%	2.2%	2.6%	3.9%	0.0%	1.1%	1.3%	1.2%	2.2%	1.7%	1.9%	2.7%			2.3%	4.6%	6.5%	7.0%	5.5%	5.8%	7.0%	7.8%	0.0%	4.5%	6.3%	6.2%	6.9%	6.6%	8.7%	9.4%			17.4%	16.0%	17.0%	18.8%	12.3%	14.1%	17.7%	19.0%	7.7%	10.6%	14.9%	17.4%	14.0%	16.2%	16.7%	19.1%			11.8%	29.3%	26.4%	34.4%	20.3%	26.0%	27.2%	30.5%	30.6%	26.8%	25.7%	23.4%	24.0%	27.3%	28.7%	31.2%			39.6%	33.3%	27.3%	36.5%	28.4%	42.0%	33.3%	34.8%	43.2%	41.2%	35.5%	36.2%	36.4%	33.2%	35.0%	33.2%			0.0%	5.5%	3.0%	2.1%	3.2%	2.2%	3.8%	2.3%	1.4%	1.8%	1.4%	1.6%	2.0%	2.6%	4.2%	4.0%			11.8%	6.0%	8.5%	9.1%	5.1%	7.7%	9.5%	9.7%	5.0%	5.0%	3.4%	9.4%	4.5%	6.6%	7.3%	9.8%			8.5%	9.6%	15.3%	18.5%	10.0%	13.5%	17.2%	20.6%	11.1%	10.5%	12.6%	16.7%	10.6%	15.7%	18.2%	21.1%			28.1%	34.5%	27.4%	32.3%	17.9%	21.0%	27.0%	29.2%	22.7%	21.9%	30.0%	24.2%	20.8%	28.3%	29.3%	33.5%			21.6%	30.2%	33.7%	36.8%	25.7%	32.7%	33.0%	34.2%	27.8%	25.5%	32.5%	33.5%	28.9%	34.4%	36.9%	39.9%			שיעור בעלי יתום HbA1c מאוזנת בחולי סוכרת ו SMI בני 18-84																	18-44	45-64	65+	18-44	45-64	65+	18-44	45-64	65+	NA*	NA*	NA*	NA*	83.7%	67.7%	74.3%	70.7%	NA*	NA*	59.4%	NA*	76.7%	73.5%	72.4%	70.6%	18-44	NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																				
0.8%	0.8%	0.9%	0.9%	1.3%	1.6%	1.9%	2.1%	1.9%	1.9%	2.1%	2.7%	1.1%	1.4%	1.7%	1.6%			שיעור ההימצאות של סוכרת בקרב חולי SMI בני 18 ומעלה																	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	1.7%	2.5%	1.3%	1.8%	2.0%	2.2%	2.6%	3.9%	0.0%	1.1%	1.3%	1.2%	2.2%	1.7%	1.9%	2.7%			2.3%	4.6%	6.5%	7.0%	5.5%	5.8%	7.0%	7.8%	0.0%	4.5%	6.3%	6.2%	6.9%	6.6%	8.7%	9.4%			17.4%	16.0%	17.0%	18.8%	12.3%	14.1%	17.7%	19.0%	7.7%	10.6%	14.9%	17.4%	14.0%	16.2%	16.7%	19.1%			11.8%	29.3%	26.4%	34.4%	20.3%	26.0%	27.2%	30.5%	30.6%	26.8%	25.7%	23.4%	24.0%	27.3%	28.7%	31.2%			39.6%	33.3%	27.3%	36.5%	28.4%	42.0%	33.3%	34.8%	43.2%	41.2%	35.5%	36.2%	36.4%	33.2%	35.0%	33.2%			0.0%	5.5%	3.0%	2.1%	3.2%	2.2%	3.8%	2.3%	1.4%	1.8%	1.4%	1.6%	2.0%	2.6%	4.2%	4.0%			11.8%	6.0%	8.5%	9.1%	5.1%	7.7%	9.5%	9.7%	5.0%	5.0%	3.4%	9.4%	4.5%	6.6%	7.3%	9.8%			8.5%	9.6%	15.3%	18.5%	10.0%	13.5%	17.2%	20.6%	11.1%	10.5%	12.6%	16.7%	10.6%	15.7%	18.2%	21.1%			28.1%	34.5%	27.4%	32.3%	17.9%	21.0%	27.0%	29.2%	22.7%	21.9%	30.0%	24.2%	20.8%	28.3%	29.3%	33.5%			21.6%	30.2%	33.7%	36.8%	25.7%	32.7%	33.0%	34.2%	27.8%	25.5%	32.5%	33.5%	28.9%	34.4%	36.9%	39.9%			שיעור בעלי יתום HbA1c מאוזנת בחולי סוכרת ו SMI בני 18-84																	18-44	45-64	65+	18-44	45-64	65+	18-44	45-64	65+	NA*	NA*	NA*	NA*	83.7%	67.7%	74.3%	70.7%	NA*	NA*	59.4%	NA*	76.7%	73.5%	72.4%	70.6%	18-44	NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																						
שיעור ההימצאות של סוכרת בקרב חולי SMI בני 18 ומעלה																	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+	18-34	35-44	45-54	55-64	65+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																												
1.7%	2.5%	1.3%	1.8%	2.0%	2.2%	2.6%	3.9%	0.0%	1.1%	1.3%	1.2%	2.2%	1.7%	1.9%	2.7%			2.3%	4.6%	6.5%	7.0%	5.5%	5.8%	7.0%	7.8%	0.0%	4.5%	6.3%	6.2%	6.9%	6.6%	8.7%	9.4%			17.4%	16.0%	17.0%	18.8%	12.3%	14.1%	17.7%	19.0%	7.7%	10.6%	14.9%	17.4%	14.0%	16.2%	16.7%	19.1%			11.8%	29.3%	26.4%	34.4%	20.3%	26.0%	27.2%	30.5%	30.6%	26.8%	25.7%	23.4%	24.0%	27.3%	28.7%	31.2%			39.6%	33.3%	27.3%	36.5%	28.4%	42.0%	33.3%	34.8%	43.2%	41.2%	35.5%	36.2%	36.4%	33.2%	35.0%	33.2%			0.0%	5.5%	3.0%	2.1%	3.2%	2.2%	3.8%	2.3%	1.4%	1.8%	1.4%	1.6%	2.0%	2.6%	4.2%	4.0%			11.8%	6.0%	8.5%	9.1%	5.1%	7.7%	9.5%	9.7%	5.0%	5.0%	3.4%	9.4%	4.5%	6.6%	7.3%	9.8%			8.5%	9.6%	15.3%	18.5%	10.0%	13.5%	17.2%	20.6%	11.1%	10.5%	12.6%	16.7%	10.6%	15.7%	18.2%	21.1%			28.1%	34.5%	27.4%	32.3%	17.9%	21.0%	27.0%	29.2%	22.7%	21.9%	30.0%	24.2%	20.8%	28.3%	29.3%	33.5%			21.6%	30.2%	33.7%	36.8%	25.7%	32.7%	33.0%	34.2%	27.8%	25.5%	32.5%	33.5%	28.9%	34.4%	36.9%	39.9%			שיעור בעלי יתום HbA1c מאוזנת בחולי סוכרת ו SMI בני 18-84																	18-44	45-64	65+	18-44	45-64	65+	18-44	45-64	65+	NA*	NA*	NA*	NA*	83.7%	67.7%	74.3%	70.7%	NA*	NA*	59.4%	NA*	76.7%	73.5%	72.4%	70.6%	18-44	NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																								
2.3%	4.6%	6.5%	7.0%	5.5%	5.8%	7.0%	7.8%	0.0%	4.5%	6.3%	6.2%	6.9%	6.6%	8.7%	9.4%			17.4%	16.0%	17.0%	18.8%	12.3%	14.1%	17.7%	19.0%	7.7%	10.6%	14.9%	17.4%	14.0%	16.2%	16.7%	19.1%			11.8%	29.3%	26.4%	34.4%	20.3%	26.0%	27.2%	30.5%	30.6%	26.8%	25.7%	23.4%	24.0%	27.3%	28.7%	31.2%			39.6%	33.3%	27.3%	36.5%	28.4%	42.0%	33.3%	34.8%	43.2%	41.2%	35.5%	36.2%	36.4%	33.2%	35.0%	33.2%			0.0%	5.5%	3.0%	2.1%	3.2%	2.2%	3.8%	2.3%	1.4%	1.8%	1.4%	1.6%	2.0%	2.6%	4.2%	4.0%			11.8%	6.0%	8.5%	9.1%	5.1%	7.7%	9.5%	9.7%	5.0%	5.0%	3.4%	9.4%	4.5%	6.6%	7.3%	9.8%			8.5%	9.6%	15.3%	18.5%	10.0%	13.5%	17.2%	20.6%	11.1%	10.5%	12.6%	16.7%	10.6%	15.7%	18.2%	21.1%			28.1%	34.5%	27.4%	32.3%	17.9%	21.0%	27.0%	29.2%	22.7%	21.9%	30.0%	24.2%	20.8%	28.3%	29.3%	33.5%			21.6%	30.2%	33.7%	36.8%	25.7%	32.7%	33.0%	34.2%	27.8%	25.5%	32.5%	33.5%	28.9%	34.4%	36.9%	39.9%			שיעור בעלי יתום HbA1c מאוזנת בחולי סוכרת ו SMI בני 18-84																	18-44	45-64	65+	18-44	45-64	65+	18-44	45-64	65+	NA*	NA*	NA*	NA*	83.7%	67.7%	74.3%	70.7%	NA*	NA*	59.4%	NA*	76.7%	73.5%	72.4%	70.6%	18-44	NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																										
17.4%	16.0%	17.0%	18.8%	12.3%	14.1%	17.7%	19.0%	7.7%	10.6%	14.9%	17.4%	14.0%	16.2%	16.7%	19.1%			11.8%	29.3%	26.4%	34.4%	20.3%	26.0%	27.2%	30.5%	30.6%	26.8%	25.7%	23.4%	24.0%	27.3%	28.7%	31.2%			39.6%	33.3%	27.3%	36.5%	28.4%	42.0%	33.3%	34.8%	43.2%	41.2%	35.5%	36.2%	36.4%	33.2%	35.0%	33.2%			0.0%	5.5%	3.0%	2.1%	3.2%	2.2%	3.8%	2.3%	1.4%	1.8%	1.4%	1.6%	2.0%	2.6%	4.2%	4.0%			11.8%	6.0%	8.5%	9.1%	5.1%	7.7%	9.5%	9.7%	5.0%	5.0%	3.4%	9.4%	4.5%	6.6%	7.3%	9.8%			8.5%	9.6%	15.3%	18.5%	10.0%	13.5%	17.2%	20.6%	11.1%	10.5%	12.6%	16.7%	10.6%	15.7%	18.2%	21.1%			28.1%	34.5%	27.4%	32.3%	17.9%	21.0%	27.0%	29.2%	22.7%	21.9%	30.0%	24.2%	20.8%	28.3%	29.3%	33.5%			21.6%	30.2%	33.7%	36.8%	25.7%	32.7%	33.0%	34.2%	27.8%	25.5%	32.5%	33.5%	28.9%	34.4%	36.9%	39.9%			שיעור בעלי יתום HbA1c מאוזנת בחולי סוכרת ו SMI בני 18-84																	18-44	45-64	65+	18-44	45-64	65+	18-44	45-64	65+	NA*	NA*	NA*	NA*	83.7%	67.7%	74.3%	70.7%	NA*	NA*	59.4%	NA*	76.7%	73.5%	72.4%	70.6%	18-44	NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																												
11.8%	29.3%	26.4%	34.4%	20.3%	26.0%	27.2%	30.5%	30.6%	26.8%	25.7%	23.4%	24.0%	27.3%	28.7%	31.2%			39.6%	33.3%	27.3%	36.5%	28.4%	42.0%	33.3%	34.8%	43.2%	41.2%	35.5%	36.2%	36.4%	33.2%	35.0%	33.2%			0.0%	5.5%	3.0%	2.1%	3.2%	2.2%	3.8%	2.3%	1.4%	1.8%	1.4%	1.6%	2.0%	2.6%	4.2%	4.0%			11.8%	6.0%	8.5%	9.1%	5.1%	7.7%	9.5%	9.7%	5.0%	5.0%	3.4%	9.4%	4.5%	6.6%	7.3%	9.8%			8.5%	9.6%	15.3%	18.5%	10.0%	13.5%	17.2%	20.6%	11.1%	10.5%	12.6%	16.7%	10.6%	15.7%	18.2%	21.1%			28.1%	34.5%	27.4%	32.3%	17.9%	21.0%	27.0%	29.2%	22.7%	21.9%	30.0%	24.2%	20.8%	28.3%	29.3%	33.5%			21.6%	30.2%	33.7%	36.8%	25.7%	32.7%	33.0%	34.2%	27.8%	25.5%	32.5%	33.5%	28.9%	34.4%	36.9%	39.9%			שיעור בעלי יתום HbA1c מאוזנת בחולי סוכרת ו SMI בני 18-84																	18-44	45-64	65+	18-44	45-64	65+	18-44	45-64	65+	NA*	NA*	NA*	NA*	83.7%	67.7%	74.3%	70.7%	NA*	NA*	59.4%	NA*	76.7%	73.5%	72.4%	70.6%	18-44	NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																														
39.6%	33.3%	27.3%	36.5%	28.4%	42.0%	33.3%	34.8%	43.2%	41.2%	35.5%	36.2%	36.4%	33.2%	35.0%	33.2%			0.0%	5.5%	3.0%	2.1%	3.2%	2.2%	3.8%	2.3%	1.4%	1.8%	1.4%	1.6%	2.0%	2.6%	4.2%	4.0%			11.8%	6.0%	8.5%	9.1%	5.1%	7.7%	9.5%	9.7%	5.0%	5.0%	3.4%	9.4%	4.5%	6.6%	7.3%	9.8%			8.5%	9.6%	15.3%	18.5%	10.0%	13.5%	17.2%	20.6%	11.1%	10.5%	12.6%	16.7%	10.6%	15.7%	18.2%	21.1%			28.1%	34.5%	27.4%	32.3%	17.9%	21.0%	27.0%	29.2%	22.7%	21.9%	30.0%	24.2%	20.8%	28.3%	29.3%	33.5%			21.6%	30.2%	33.7%	36.8%	25.7%	32.7%	33.0%	34.2%	27.8%	25.5%	32.5%	33.5%	28.9%	34.4%	36.9%	39.9%			שיעור בעלי יתום HbA1c מאוזנת בחולי סוכרת ו SMI בני 18-84																	18-44	45-64	65+	18-44	45-64	65+	18-44	45-64	65+	NA*	NA*	NA*	NA*	83.7%	67.7%	74.3%	70.7%	NA*	NA*	59.4%	NA*	76.7%	73.5%	72.4%	70.6%	18-44	NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																
0.0%	5.5%	3.0%	2.1%	3.2%	2.2%	3.8%	2.3%	1.4%	1.8%	1.4%	1.6%	2.0%	2.6%	4.2%	4.0%			11.8%	6.0%	8.5%	9.1%	5.1%	7.7%	9.5%	9.7%	5.0%	5.0%	3.4%	9.4%	4.5%	6.6%	7.3%	9.8%			8.5%	9.6%	15.3%	18.5%	10.0%	13.5%	17.2%	20.6%	11.1%	10.5%	12.6%	16.7%	10.6%	15.7%	18.2%	21.1%			28.1%	34.5%	27.4%	32.3%	17.9%	21.0%	27.0%	29.2%	22.7%	21.9%	30.0%	24.2%	20.8%	28.3%	29.3%	33.5%			21.6%	30.2%	33.7%	36.8%	25.7%	32.7%	33.0%	34.2%	27.8%	25.5%	32.5%	33.5%	28.9%	34.4%	36.9%	39.9%			שיעור בעלי יתום HbA1c מאוזנת בחולי סוכרת ו SMI בני 18-84																	18-44	45-64	65+	18-44	45-64	65+	18-44	45-64	65+	NA*	NA*	NA*	NA*	83.7%	67.7%	74.3%	70.7%	NA*	NA*	59.4%	NA*	76.7%	73.5%	72.4%	70.6%	18-44	NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																		
11.8%	6.0%	8.5%	9.1%	5.1%	7.7%	9.5%	9.7%	5.0%	5.0%	3.4%	9.4%	4.5%	6.6%	7.3%	9.8%			8.5%	9.6%	15.3%	18.5%	10.0%	13.5%	17.2%	20.6%	11.1%	10.5%	12.6%	16.7%	10.6%	15.7%	18.2%	21.1%			28.1%	34.5%	27.4%	32.3%	17.9%	21.0%	27.0%	29.2%	22.7%	21.9%	30.0%	24.2%	20.8%	28.3%	29.3%	33.5%			21.6%	30.2%	33.7%	36.8%	25.7%	32.7%	33.0%	34.2%	27.8%	25.5%	32.5%	33.5%	28.9%	34.4%	36.9%	39.9%			שיעור בעלי יתום HbA1c מאוזנת בחולי סוכרת ו SMI בני 18-84																	18-44	45-64	65+	18-44	45-64	65+	18-44	45-64	65+	NA*	NA*	NA*	NA*	83.7%	67.7%	74.3%	70.7%	NA*	NA*	59.4%	NA*	76.7%	73.5%	72.4%	70.6%	18-44	NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																				
8.5%	9.6%	15.3%	18.5%	10.0%	13.5%	17.2%	20.6%	11.1%	10.5%	12.6%	16.7%	10.6%	15.7%	18.2%	21.1%			28.1%	34.5%	27.4%	32.3%	17.9%	21.0%	27.0%	29.2%	22.7%	21.9%	30.0%	24.2%	20.8%	28.3%	29.3%	33.5%			21.6%	30.2%	33.7%	36.8%	25.7%	32.7%	33.0%	34.2%	27.8%	25.5%	32.5%	33.5%	28.9%	34.4%	36.9%	39.9%			שיעור בעלי יתום HbA1c מאוזנת בחולי סוכרת ו SMI בני 18-84																	18-44	45-64	65+	18-44	45-64	65+	18-44	45-64	65+	NA*	NA*	NA*	NA*	83.7%	67.7%	74.3%	70.7%	NA*	NA*	59.4%	NA*	76.7%	73.5%	72.4%	70.6%	18-44	NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																						
28.1%	34.5%	27.4%	32.3%	17.9%	21.0%	27.0%	29.2%	22.7%	21.9%	30.0%	24.2%	20.8%	28.3%	29.3%	33.5%			21.6%	30.2%	33.7%	36.8%	25.7%	32.7%	33.0%	34.2%	27.8%	25.5%	32.5%	33.5%	28.9%	34.4%	36.9%	39.9%			שיעור בעלי יתום HbA1c מאוזנת בחולי סוכרת ו SMI בני 18-84																	18-44	45-64	65+	18-44	45-64	65+	18-44	45-64	65+	NA*	NA*	NA*	NA*	83.7%	67.7%	74.3%	70.7%	NA*	NA*	59.4%	NA*	76.7%	73.5%	72.4%	70.6%	18-44	NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																								
21.6%	30.2%	33.7%	36.8%	25.7%	32.7%	33.0%	34.2%	27.8%	25.5%	32.5%	33.5%	28.9%	34.4%	36.9%	39.9%			שיעור בעלי יתום HbA1c מאוזנת בחולי סוכרת ו SMI בני 18-84																	18-44	45-64	65+	18-44	45-64	65+	18-44	45-64	65+	NA*	NA*	NA*	NA*	83.7%	67.7%	74.3%	70.7%	NA*	NA*	59.4%	NA*	76.7%	73.5%	72.4%	70.6%	18-44	NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																										
שיעור בעלי יתום HbA1c מאוזנת בחולי סוכרת ו SMI בני 18-84																	18-44	45-64	65+	18-44	45-64	65+	18-44	45-64	65+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																		
NA*	NA*	NA*	NA*	83.7%	67.7%	74.3%	70.7%	NA*	NA*	59.4%	NA*	76.7%	73.5%	72.4%	70.6%	18-44	NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																						
NA*	70.9%	68.9%	61.0%	74.7%	76.3%	68.3%	68.2%	NA*	76.9%	68.5%	59.4%	75.0%	73.9%	74.1%	69.5%	45-64	NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
NA*	87.1%	91.4%	NA*	89.1%	83.3%	79.5%	76.2%	84.4%	84.5%	86.9%	82.9%	86.9%	83.6%	83.4%	74.5%	65+	NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																								
NA*	NA*	NA*	NA*	97.2%	79.2%	78.4%	NA*	NA*	NA*	NA*	NA*	86.7%	75.6%	64.9%	70.0%	18-44	NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																									
NA*	76.9%	79.2%	71.2%	87.7%	81.3%	76.2%	77.6%	NA*	80.7%	78.3%	63.4%	80.3%	77.4%	76.1%	69.4%	45-64	NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																										
NA*	76.6%	87.2%	NA*	85.3%	87.5%	81.9%	85.9%	NA*	83.9%	73.4%	87.0%	90.6%	84.0%	80.9%	82.9%	65+	שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																											
שיעור היתעד של מרכיבי BMI בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																		
50.0%	53.3%	46.0%	41.2%	28.6%	29.8%	29.2%	28.7%	24.2%	30.0%	28.2%	24.3%	37.4%	36.2%	36.7%	32.9%	20-34	45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																						
45.3%	52.0%	49.4%	49.1%	32.7%	30.1%	35.0%	33.1%	28.0%	35.2%	32.9%	35.4%	44.3%	43.4%	46.1%	46.0%	35-44	62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
62.8%	59.1%	61.3%	63.8%	39.4%	45.4%	46.0%	47.8%	46.0%	50.6%	47.3%	46.8%	59.2%	63.1%	63.2%	64.2%	45-64	55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																								
55.3%	54.5%	64.8%	64.7%	35.5%	40.0%	43.2%	51.6%	37.3%	41.9%	45.6%	48.3%	41.9%	46.1%	49.3%	44.9%	20-34	49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																									
49.0%	55.5%	67.9%	68.5%	38.2%	41.8%	47.8%	42.9%	50.8%	51.4%	43.9%	54.5%	45.9%	53.5%	54.1%	54.0%	35-44	61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																										
61.9%	69.4%	68.8%	74.3%	43.7%	50.3%	52.4%	55.9%	49.7%	58.3%	57.8%	59.1%	64.2%	68.9%	69.1%	70.9%	45-64	שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64	NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																											
שיעור ההימצאות של השמנת יתר בקרב חולי SMI בני 20-64																	20-34	35-44	45-64	20-34	35-44	45-64	20-34	35-44	45-64																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																		
NA*	33.7%	36.3%	35.7%	33.2%	39.3%	38.4%	40.4%	32.3%	40.4%	32.8%	38.1%	35.0%	35.6%	37.8%	38.1%	20-34	23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																						
23.1%	42.3%	48.4%	53.4%	40.0%	40.8%	47.2%	49.4%	NA*	37.5%	42.6%	39.7%	37.0%	40.9%	42.0%	38.5%	35-44	37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
37.4%	41.2%	39.3%	45.5%	38.8%	41.3%	45.6%	48.3%	32.0%	40.9%	40.1%	41.4%	35.7%	36.2%	36.4%	36.8%	45-64	NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																								
NA*	36.7%	56.6%	55.7%	36.2%	35.5%	47.8%	47.9%	NA*	38.9%	43.6%	50.4%	35.9%	42.5%	44.9%	51.0%	20-34	NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																									
NA*	51.5%	54.9%	59.0%	36.1%	45.7%	49.3%	56.4%	38.7%	47.3%	54.5%	61.4%	37.0%	46.6%	54.9%	56.5%	35-44	36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																										
36.0%	53.8%	52.4%	68.5%	41.0%	49.1%	53.9%	58.8%	48.6%	44.1%	54.2%	59.4%	41.5%	47.1%	51.5%	56.4%	45-64	שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+	NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																											
שיעור המפגשים הטפילים עם אנשי בריאות הנפש בקהילה תוך 14 יום משחרור מאשפוז פסיכיאטרי ממושך																	18-34	35-54	55+	18-34	35-54	55+	18-34	35-54	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																		
NA*	38.8%	26.4%	30.9%	39.2%	45.1%	32.5%	33.8%	NA*	25.0%	11.2%	16.5%	87.2%	93.2%	90.2%	91.3%	18-34	NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																						
NA*	44.9%	27.8%	25.0%	29.6%	36.2%	41.7%	45.6%	NA*	7.0%	11.5%	2.5%	89.1%	89.9%	90.8%	93.4%	35-54	NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																							
NA*	NA*	NA*	NA*	35.7%	28.6%	25.0%	44.1%	NA*	NA*	12.5%	NA*	93.5%	95.5%	87.7%	87.5%	55+	NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																								
NA*	45.0%	40.4%	46.8%	55.7%	37.8%	39.1%	37.5%	NA*	NA*	25.8%	11.8%	89.4%	89.7%	91.4%	89.3%	18-34	NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																									
NA*	33.3%	40.0%	NA*	45.5%	47.2%	39.1%	31.0%	NA*	NA*	14.6%	NA*	96.3%	92.9%	92.4%	91.1%	35-54	NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																										
NA*	NA*	39.4%	NA*	40.5%	43.0%	32.3%	25.0%	NA*	NA*	NA*	NA*	95.0%	91.0%	90.7%	92.2%	55+																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																											

תא מסומן ב-* מייצג ערך מכנה נומן מ 30 מבטוחים